arXiv:0912.2715v3 [math.GT] 30 Oct 2011

A COMBINATION THEOREM FOR METRIC BUNDLES

MAHAN MJ AND PRANAB SARDAR

Abstract. We define metric bundles/metric graph bundles which provide a purely topological/coarse-geometric generalization of the notion of trees of metric spaces a la Bestvina-Feighn in the special case that the inclusions of the edge spaces into the vertex spaces are uniform coarsely surjective quasi-isometries. We prove the existence of quasi-isometric sections in this generality. Then we prove a combination theorem for metric (graph) bundles (including exact sequences of groups) that establishes sufficient conditions, particularly flaring, under which the metric bundles are hyperbolic. We use this to give examples of surface bundles over hyperbolic disks, whose universal cover is Gromov-hyperbolic. We also show that in typical situations, flaring is also a necessary condition.

Contents

1. Introduction 2
1.1. Metric Bundles 5
1.2. Hyperbolic metric spaces 10
1.3. Trees of hyperbolic and relatively hyperbolic metric spaces 14
2. QI Sections 19
2.1. Existence of qi sections 19
2.2. Nearby qi sections 27
3. Construction of Hyperbolic Ladders 29
3.1. Hyperbolicity of ladders: Special case 31
3.2. Hyperbolicity of ladders: General case 37
4. The Combination Theorem 41
4.1. Proof of Main Theorem 41
5. Consequences and Applications 45
5.1. Sections, Retracts and Cannon-Thurston maps 46
5.2. Hyperbolicity of base and flaring 47
5.3. Necessity of Flaring 49
5.4. An Example 52
References 55

Date: November 1, 2011.

Research of first author partially supported by a CEFIPRA Indo-French Research grant. The second author is partly supported by a CSIR Research Fellowship. This paper is part of PS's PhD thesis written under the supervision of MM. .

1. Introduction

In this paper we introduce the notions of metric bundles (resp. metric graph bundles) which provide a purely topological (resp. coarse-geometric) generalization of the notion of trees of metric spaces a la Bestvina-Feighn [BF92] in the special case that the inclusions of the edge spaces into the vertex spaces are uniform coarsely surjective quasi-isometries. We generalize the base space from a tree to an arbitrary hyperbolic metric space. In [FM02], Farb and Mosher introduced the notion of metric fibrations which was used by Hamenstadt to give a combination theorem in [Ham05]. Metric fibrations can be thought of as metric bundles (in our terminology) equipped with a *foliation by totally geodesic sections* of the base space. We first prove the following Proposition which ensures the existence of *q(uasi)-i(sometric)* sections in the general context of metric bundles, generalizing and giving a different proof of a Lemma of Mosher [Mos96] in the context of exact sequences of groups (see Example 1.5).

Proposition 2.10 (Existence of qi sections): *Let* $\delta, N \geq 0$ *and suppose* $p : X \to B$ *is an* (f, K)*-metric graph bundle with the following properties:*

1. *Each of the fibers* F_b*,* $b \in \mathcal{V}(B)$ *is a* δ*-hyperbolic geodesic metric space with respect to the path metric* d_b *induced from* X*.*
2. *The barycenter maps* $\phi_b : \partial^3 F_b \to F_b$ *are uniformly coarsely surjective, i.e. the image of the map* ϕ_b *is contained in the vertex set* $\mathcal{V}(F_b)$ *and* F_b *is contained in the* N*-neighborhood of the image of* ϕ_b *for all* $b \in \mathcal{V}(B)$*.*

Then there is a $K_0 = K_0(f, K, \delta, N)$*-qi section through each point of* $\mathcal{V}(X)$*.*

Proposition 2.10 provides a context for developing a 'coarse theory of bundles' and proving the following combination theorem, which is the main theorem of this paper.

Theorem 4.10: *Suppose* $p : X \to B$ *is a metric bundle (resp. metric graph bundle) such that*
(1) B *is a* δ*-hyperbolic metric space.*
(2) *Each of the fibers* F_b*,* $b \in B$ *(resp.* $b \in \mathcal{V}(B)$*) is a* δ'*-hyperbolic geodesic metric space with respect to the path metric induced from* X*.*
(3) *The barycenter maps* $\partial^3 F_b \to F_b$*,* $b \in B$ *(resp.* $b \in \mathcal{V}(B)$*) are uniformly coarsely surjective.*
(4) *A flaring condition is satisfied.*
Then X *is a hyperbolic metric space.*

This is a first step towards proving a combination Theorem for more general complexes of spaces (cf. Problem 90 of [Kap08], part of the Geometric Group Theory problem list).

Theorem 4.10 generalizes Hamenstadt's combination theorem (Corollary 3.8 of [Ham05]) in two ways:
a) It removes the hypothesis of properness of the base space B – a hypothesis that is crucial in [Ham05] to ensure compactness of the boundary of the base space and hence allow the arguments in [Ham05] to work. This generalization is relevant for two reasons. First, underlying trees in trees of spaces are frequently non-proper. Secondly, curve complexes of surfaces are mostly non-proper metric spaces and occur as natural base spaces for metric bundles. See [LMS08] by Leininger-Mj-Schleimer for a closely related example.

b) It removes the hypothesis on existence of totally geodesic sections in [Ham05] altogether. Proposition 2.10 ensures the existence of qi sections under mild technical assumptions.

A word about the proof ahead of time. Proposition 2.10 ensures the existence of qi sections through points of X. We use the notion of flaring from Bestvina-Feighn [BF92] and a criterion for hyperbolicity introduced by Hamenstadt in [Ham07] to construct certain path families and use them to prove hyperbolicity. Another crucial ingredient is a 'ladder-construction' due to Mitra [Mit98b], which may be regarded as an analog of the hallways of [BF92].

Recall [Far98] that for a pair $(X, \mathcal{H})$ of a metric space (X, d_X) and a family of path-connected subsets $\mathcal{H}$ of X, the electric space $\mathcal{E}(X, \mathcal{H})$ is the pseudo-metric space $X \sqcup_{H\in\mathcal{H}} H \times [0,1]$ with $H \times \{0\}$ identified with $H \subset X$ and $H \times \{1\}$ equipped with the zero metric. Each $\{h\} \times [0,1]$ is isometric to the unit interval. There is a natural inclusion map $E : X \to \mathcal{E}(X, \mathcal{H})$ which is referred to as the electrocution map. The image $E(X)$ inherits a metric called the electric metric d_e.

As an application of Theorem 4.10 we obtain a rather plentiful supply of examples from the following Proposition, where the base space need not be a tree (as in all previously known examples). Let S be a closed surface of genus greater than one and $Teich(S)$ be the Teichmuller space of S. The Teichmuller metric on $Teich(S)$ is denoted as d_T and d_e denotes the electric metric on $Teich(S)$ obtained by electrocuting the cusps of the moduli space $Mod(S)$ and lifting the resulting metric to $Teich(S)$.

Proposition 5.15: *Let (K, d_K) be a hyperbolic metric space satisfying the following:*
There exists $C \geq 0$ such that for any two points $u, v \in K$, there exists a bi-infinite C-quasigeodesic $\gamma \subset K$ with $d_K(u,\gamma) \leq C$ and $d_K(v,\gamma) \leq C$.
Let $j : K \to (Teich(S), d_T)$ be a quasi-isometric embedding such that $E \circ j : K \to (Teich(S), d_e)$ is also a quasi-isometric embedding. Let $U(S,K)$ be the pull-back bundle under j of the universal curve over $Teich(S)$ equipped with the natural path metric. Then $\widetilde{U(S,K)}$ is a hyperbolic metric space, where $\widetilde{U(S,K)}$ denotes the metric bundle obtained by taking fiberwise universal covers.

It is an open question (cf. [KL08] [FM02]) to find purely pseudo Anosov surface groups Q $(= \pi_1(\Sigma)$, say) in $MCG(S)$. This is equivalent to constructing surface bundles over surfaces with total space W, fiber S, and base Σ, such that $\pi_1(W)$ does not contain a copy of $\mathbb{Z}\oplus\mathbb{Z}$. One way of ensuring this is to find an example where the total space has (Gromov) hyperbolic fundamental group $\pi_1(W)$. A quasi-isometric model for the universal cover $\widetilde{W}$ is a metric graph bundle where the fibers are Cayley graphs of $\pi_1(S)$ and the base K a Cayley graph of $\pi_1(\Sigma)$. Using a construction of Leininger and Schleimer [LS11] in conjunction with Proposition 5.15 we construct examples of hyperbolic metric graph bundles where fibers are Cayley graphs of $\pi_1(S)$ and K is a hyperbolic disk. However the disks K are not invariant under a surface group; so we only obtain surface bundles W over K with fiber S such that the universal cover $\widetilde{W}$ is hyperbolic.

We also obtain the following characterization of convex cocompact subgroups of mapping class groups of surfaces S^h with punctures. We state the result for surfaces with a single puncture.

Proposition 5.17: *Let* $K = \pi_1(S^h)$ *be the fundamental group of a surface with a single puncture and let* K_1 *be its peripheral subgroup. Let* Q *be a convex cocompact subgroup of the mapping class group of* S^h*. Let*

$$1 \to (K, K_1) \to (G, N_G(K_1)) \xrightarrow{p} (Q, Q_1) \to 1$$

be the induced short exact sequence of (pairs of) groups with $Q_1 = N_G(K_1)/K_1$*. Then* G *is (strongly) hyperbolic relative to* $N_G(K_1)$*.*

Conversely, if G *is (strongly) hyperbolic relative to* $N_G(K_1)$*, then* Q *is convex-cocompact.*

Theorem 4.10 also provides the following combination theorem whenever we have an exact sequence with hyperbolic quotient and kernel. This gives a converse to a theorem of Mosher [Mos96].

Theorem 5.1: *Suppose that the short exact sequence of finitely generated groups*

$$1 \to K \to G \to Q \to 1$$

satisfies a flaring condition such that K, Q *are word hyperbolic and* K *is non-elementary. Then* G *is hyperbolic.*

The next Proposition links the flaring condition to hyperbolicity of the base.

Proposition 5.5: *Consider the short exact sequence of finitely generated groups*

$$1 \to K \to G \to Q \to 1$$

such that K *is non-elementary word hyperbolic but* Q *is not hyperbolic. Then the short exact sequence cannot satisfy a flaring condition.*

We also prove an analog of Proposition 5.5 for relatively hyperbolic groups and use it to generalize a Theorem of Mosher [Mos96] as follows.

Proposition 5.7: *Suppose we have a short exact sequence of finitely generated groups*

$$1 \to (K, K_1) \to (G, N_G(K_1)) \xrightarrow{p} (Q, Q_1) \to 1$$

with K *(strongly) hyperbolic relative to the cusp subgroup* K_1 *such that* G *preserves cusps and* $Q_1 = N_G(K_1)/K_1$*. Suppose further that* G *is (strongly) hyperbolic relative to* $N_G(K_1)$*. Then* $Q(= Q_1)$ *is hyperbolic.*

Finally we show the necessity of flaring.

Proposition 5.8: *Let* $P : X \to B$ *be a metric (graph) bundle such that*
1) X *is hyperbolic.*
2) There exists δ_0 *such that each fiber* $F_z = p^{-1}(z) \subset X$ *equipped with the inherited path metric is* δ_0*-hyperbolic.*
Then the metric bundle satisfies a flaring condition.

In particular, any exact sequence of finitely generated groups $1 \to N \to G \to Q \to 1$ *with* N *and* G *hyperbolic, satisfies a flaring condition.*

Outline of the main steps:
There are four main steps in the proof of Combination Theorem 4.10. Precise definitions of terms are given in the next subsection.
1) First we construct a metric graph bundle (see Definition 1.3) out of a given metric bundle. The bundles have quasi-isometric base space and total space. Next we set out to prove that this metric graph bundle is hyperbolic under the given conditions on the metric bundle.
2) Proposition 2.10 proves the existence of qi sections and is the coarse geometric analog of the statement that a principal bundle with contractible base is trivial.

The main ingredient of the proof is the definition of a 'discrete flow' of one fiber to another fiber. This is the content of Section 2.1. The main idea is elaborated upon in the first paragraph of Section 2.1.
3) Any two such qi sections bound a 'ladder' between them (cf. Definition 3.1, [Mit98a], [Mit98b]). The next step is to prove the hyperbolicity of these ladders. In Section 3.1 we prove hyperbolicity of small-girth ladders (Proposition 3.6). In Section 3.2 we break up a big ladder into small-girth ladders and use a consequence (Proposition 1.41) of a combination theorem due to Mj-Reeves [MR08] to conclude that the whole ladder is hyperbolic.
4) In Section 4, we assemble the pieces to prove Theorem 4.10.

For the reader interested in getting to the main ideas of Theorem 4.10 without getting into technical details, we have sketched Steps (1) and (2) above in the first paragraph of Section 2.1, the first paragraph of Section 3 and the paragraph following the statement of Proposition 3.6 in Section 3.1.

Acknowledgments: We would like to thank Panos Papasoglu for explaining the proof of the last statement of Theorem 5.4 to us. We would also like to thank Chris Leininger for sharing his examples in [LS11] with us. This paper owes an intellectual debt to Hamenstadt's paper [Ham05], which inspired us to find a combination Theorem in the generality described here. Finally we would like to thank the referee for a meticulous reading of the manuscript and for several helpful remarks and comments. In particular, the notion of a metric graph bundle arose out of the referee's comments on an earlier draft.

1.1. Metric Bundles.

Let X, Y be metric spaces and let $k \geq 1$, $\epsilon \geq 0$. Recall [Gro85] [Gd90] that a map $\phi : X \to Y$ is said to be a (k, ϵ)-**quasi-isometric embedding** if $\forall x_1, x_2 \in X$ one has

$$d(x_1, x_2)/k - \epsilon \leq d(\phi(x_1), \phi(x_2)) \leq k.d(x_1, x_2) + \epsilon.$$

A (k, k)-quasi-isometric embedding will simply be referred to as a k-**quasi-isometric embedding**. A map $\phi : X \to Y$ is said to be a (k, ϵ)-**quasi-isometry** (resp. k-**quasi-isometry**) if it is a (k, ϵ)-quasi-isometric embedding (resp. k-quasi-isometric embedding) and if there is a constant $D > 0$ such that $\forall\, y \in Y$, $\exists x \in X$ with $d(\phi(x), y) \leq D$.

A (k, ϵ)-**quasi-geodesic** (resp. a k-**quasi-geodesic**) in a metric space X is a (k, ϵ)-quasi-isometric embedding (resp. a k-quasi-isometric embedding) $\gamma : I \to X$, where $I \subseteq \mathbb{R}$ is an interval.

Definition 1.1. *Suppose (X, d) and (B, d_B) are geodesic metric spaces; let $c \geq 1$ and let $f : \mathbb{R}^+ \to \mathbb{R}^+$ be a function. We say that X is an $(f, c)-$* **metric bundle** *over B if there is a surjective 1-Lipschitz map $p : X \to B$ such that the following conditions hold:*
1) For each point $z \in B$, $F_z := p^{-1}(z)$ is a geodesic metric space with respect to the path metric d_z induced from X. The inclusion maps $i : (F_z, d_z) \to X$ are uniformly metrically proper, i.e. for all $z \in B$ and $x, y \in F_z$, $d(i(x), i(y)) \leq N$ implies that $d_z(x, y) \leq f(N)$.
2) *Suppose $z_1, z_2 \in B$, $d(z_1, z_2) \leq 1$ and let γ be a geodesic in B joining them.*
$2(i)$ *Then for any point $x \in F_z$, $z \in \gamma$, there is a path in $p^{-1}(\gamma)$ of length at most c joining x to both F_{z_1} and F_{z_2}.*

A family of maps $f_\alpha : X_\alpha \to Y_\alpha$ between metric spaces is said to be *uniformly coarsely surjective* if there is a constant $D \geq 0$, such that for all α, Y_α is contained in the D-neighborhood of $f_\alpha(X_\alpha)$. The next Proposition proves that a certain condition ($2(ii)$ below) is a consequence of the above definition.

Proposition 1.2. *Let X be an $(f,c)-$ metric bundle over B. Then (following the notation of Definition 1.1)*
Condition 2(ii): *There exists $K \geq 1$, such that the following holds. Let $\phi : F_{z_1} \to F_{z_2}$, be any map such that $\forall x_1 \in F_{z_1}$ there is a path of length at most c in $p^{-1}(\gamma)$ joining x_1 to $\phi(x_1)$. Then ϕ is a K-quasi-isometry.*

Proof. Let $u, v \in F_{z_1}$ such that $d_{z_1}(u,v) \leq 1$. Then $d(\phi(u),\phi(v)) \leq 2c+1$ by the triangle inequality and hence $d_{z_2}(\phi(u),\phi(v)) \leq f(2c+1)$. A similar map $\overline{\phi} : F_{z_2} \to F_{z_1}$ may be defined interchanging the roles of z_1, z_2. Let $K = f(2c+1)$. Then ϕ is a K-quasi-isometric embedding. Also, $\overline{\phi}$ is a quasi-isometric inverse of ϕ as $d_{z_1}(\overline{\phi} \circ \phi(u), u) \leq f(2c)$ by a similar argument. Hence ϕ is a K-quasi-isometry. Note further that ϕ is uniformly coarsely surjective (independent of z_1, z_2). □

Since the constant K of Proposition 1.2 will be important, we will find it convenient to refer to an $(f,c)-$ metric bundle as an $(f,c,K)-$ **metric bundle** (with $K = f(2c+1)$) or simply a **metric bundle** when the parameters are not important.

For the rest of the paper by a **graph** we will always mean a connected metric graph all of whose edges are of length 1. For a graph X, $\mathcal{V}(X)$ will denote its vertex set. By a **path** in a graph we will always mean an edge path starting and ending at two vertices.

Definition 1.3. *Suppose X and B are graphs. Let $f : \mathbb{R}^+ \to \mathbb{R}^+$ be a function.*

We say that X is an $f-$ **metric graph bundle** *over B if there exists a surjective simplicial map $p : X \to B$ such that:*
1. *For each $b \in \mathcal{V}(B)$, $F_b := p^{-1}(b)$ is a connected subgraph of X and the inclusion maps $i : F_b \to X$ are uniformly metrically proper for the induced path metric d_b on F_b, i.e. for all $b \in B$ and $x, y \in F_b$, $d(i(x), i(y)) \leq N$ implies that $d_b(x,y) \leq f(N)$.*
$2(i)$ *Suppose $b_1, b_2 \in \mathcal{V}(B)$ are adjacent vertices. Then each vertex x_1 of F_{b_1} is connected by an edge with a vertex in F_{b_2}.*

As in Proposition 1.2, we have the following:

Proposition 1.4. Condition 2(ii): *There exists $K \geq 1$ such that the following holds (following the notation of Definition 1.3).*
Let $\phi : F_{b_1} \to F_{b_2}$ be any map such that each $x_1 \in \mathcal{V}(F_{b_1})$ is connected to $\phi(x_1) \in \mathcal{V}(F_{b_2})$ by an edge, and any point on an edge of F_{b_1} is sent to the image of one of the vertices on which the edge is incident. Then any such ϕ is a K-quasi-isometry.

Proof. First note that $d_{b_1}(u,v) \leq 1$ implies that $d_X(\phi(u),\phi(v)) \leq 4$ by the triangle inequality. Hence $d_{b_2}(\phi(u),\phi(v)) \leq f(4)$ since X is an $f-$ metric graph bundle. Taking $K = f(4)$, ϕ is a K-quasi-isometric embedding as in Proposition 1.2. Interchanging the roles of b_1, b_2 we obtain a quasi-isometric inverse $\overline{\phi}$ of ϕ. Also note that ϕ is uniformly coarsely surjective as in Proposition 1.2. □

Since K will be important We will find it convenient to refer to an $f-$metric graph bundle as an (f,K)**-metric graph bundle** with $(K = f(4))$ or simply as a **metric graph bundle** when f, K are understood.

For both metric bundles and metric graph bundles the spaces (F_z, d_z), $z \in B$, will be referred to as the **horizontal spaces** and the distance between two points in F_z will be referred to as their **horizontal distance**. A geodesic in F_z will be called a **horizontal geodesic**. The spaces X and B will be referred to as the *total space* and the *base space* respectively. By a statement of the form 'X is a metric bundle (resp. metric graph bundle)' we will mean that it is the total space of a metric bundle (resp. metric graph bundle).

A principal motivational example is the following.

Example 1.5. Suppose we have an exact sequence of finitely generated groups

$$1 \to N \overset{i}{\to} G \overset{\pi}{\to} Q \to 1.$$

This naturally gives rise to a metric graph bundle as follows. Choose a finite generating set S of G such that S contains a generating set S_1 of N. Let $X = \Gamma(G, S)$ be the Cayley graph of G with respect to the generating set S. Let $T = (\pi(S) \setminus \{1\})$ and $B := \Gamma(Q, T)$ be the Cayley graph of the group Q with respect to the generating set T. Then one easily checks that the map π naturally induces a simplicial map $\pi : X \to B$ between Cayley graphs, and hence a metric graph bundle structure on X over B. Note that in this case the horizontal spaces are all abstractly isometric to each other.

Another simple example to keep in mind is the following.

Example 1.6. Let $X = \mathbb{H}^2$ and $B = \mathbb{R}$. Identify B with a bi-infinite geodesic $\gamma \subset X$ with endpoints a, b on the ideal boundary. Through $x \in \gamma$, let F_x be the unique horocycle based at a. Define $p : X \to B$ by $p(F_x) = x$. This gives rise to a metric bundle structure on X over B. Note that each F_x, equipped with the induced path-metric, is abstractly isometric to $\mathbb{R}$.

A more interesting set of examples is furnished by Proposition 5.15 towards the end of the paper. We continue with the basic notions relevant to metric bundles.

Definition 1.7. *Let X be a metric bundle (resp. metric graph bundle) over a geodesic metric space B (resp. graph B) equipped with the map $p : X \to B$. Let $k \geq 1$. Then $X_1 \subseteq X$ is said to be a $k-$**quasi-isometric section of** B, or simply a $k-$qi section if there is a k-quasi-isometric embedding $s : B \to X$ (resp. $s : \mathcal{V}(B) \to \mathcal{V}(X)$) such that $p \circ s = I$ and $X_1 = Im(s)$. If X_1 is a k-qi section and $x \in X_1$, then we say that X_1 is a k-qi section through x. Also, $X_1 \subset X$ is said to be a qi section if it is a k-qi section for some $k \geq 1$.*

Definition 1.8. Let $\gamma : I \to B$ be a geodesic, where $I \subseteq \mathbb{R}$ is an interval. By a $k-$**qi lift** of γ in X, we mean a k-quasi isometric embedding $\tilde{\gamma} : I \to X$ such that $p \circ \tilde{\gamma} = \gamma$ (with the pro viso that for a metric graph bundle, I is of the form $[0, n]$ for some $n \in \mathbb{N}$, and the equality $p \circ \tilde{\gamma} = \gamma$ holds only at the integer points).
Suppose $X_1 \subseteq X$ is k- qi section and $\gamma : I \to B$ is a geodesic. By the lift of γ in X_1 we mean the $k-$qi lift of γ whose image is contained in X_1.

Definition 1.9. *Suppose $p : X \to B$ is a metric bundle or a metric graph bundle. We say that it satisfies a **flaring condition** if for all $k \geq 1$, there exist $M_k \geq 1$, $\lambda_k > 1$ and $n_k > 0$ such that the following holds:*
Let $\gamma : [-n_k, n_k] \to B$ be a geodesic and let $\tilde{\gamma_1}$ and $\tilde{\gamma_2}$ be two k-qi lifts of γ in X. If $d_{\gamma(0)}(\tilde{\gamma_1}(0), \tilde{\gamma_2}(0)) \geq M_k$, then we have

$$\lambda_k . d_{\gamma(0)}(\tilde{\gamma_1}(0), \tilde{\gamma_2}(0)) \leq max\{d_{\gamma(n_k)}(\tilde{\gamma_1}(n_k), \tilde{\gamma_2}(n_k)), d_{\gamma(-n_k)}(\tilde{\gamma_1}(-n_k), \tilde{\gamma_2}(-n_k))\}.$$

Lemma 1.10. *For every function $f : \mathbb{R}^+ \to \mathbb{R}^+$ and $K \geq 1$ there is a function $g : \mathbb{R}^+ \to \mathbb{R}^+$ such that the following holds:*

Suppose X is an (f, K)-metric graph bundle over B and $b_1, b_2 \in \mathcal{V}(B)$ with $d(b_1, b_2) = 1$. Let $C \geq 0$ and let $\phi : F_{b_1} \to F_{b_2}$ be any map such that $\forall x_1 \in F_{b_1}$, $d(x_1, \phi(x_1)) \leq C$. Then ϕ is a $g(C)$-quasi-isometry.

Proof. For $C = 2$ we know, from condition $2(i)$ of the definition of metric graph bundles, that such a map, say ϕ_0, exists; condition $2(ii)$ says that ϕ_0 is a K-quasi-isometry. For the general case, we note that $d(\phi_0(x), \phi(x)) \leq C + 2$, for all $x \in F_{b_1}$. Thus we have $d_{b_2}(\phi_0(x), \phi(x)) \leq f(C + 2)$, for all $x \in F_{b_1}$. We know that any map which is at a distance at most R from a K-quasi-isometry is a $(K + 2R)$-quasi-isometry. Thus choosing $g(C)$ to be $K + 2f(C + 2)$ we are through. □

Remark 1.11. The above lemma is also true for any (f, c, K)-metric bundle. Since the proof is an exact replica we omit it. The same applies to the following corollary as well. The corollary is immediate from the above lemma. With this in mind, later on we shall refer to these results in the context of metric bundles also.

Corollary 1.12. *For all $k \geq 1$ there is a function $\mu_k : \mathbb{R}^+ \to [1, \infty)$ such that the following holds:*

Suppose X is an (f, K)-metric graph bundle with base space B. Let $\gamma \subset B$ be a geodesic joining $b_1, b_2 \in B$, and let $\tilde{\gamma_1}$, $\tilde{\gamma_2}$ be two k-qi lifts of γ in X which join the pairs of points (x_1, x_2) and (y_1, y_2) respectively, so that $p(x_i) = p(y_i) = b_i$, $i = 1, 2$. If $d_B(b_1, b_2) \leq N$ then

$$d_{b_1}(x_1, y_1) \leq \mu_k(N) max\{d_{b_2}(x_2, y_2), 1\}.$$

Proof. Choose $\mu_k(N) = g(2k)^N$, where g is as in Lemma 1.10. □

In the rest of the paper, we will summarize the conclusion of the above corollary by saying that a metric bundle or a metric graph bundle satisfies a **bounded flaring condition**.

We end this subsection by showing that a metric bundle naturally gives rise to a metric graph bundle, such that the respective base and total spaces are quasi-isometric. We recall a general fact that geodesic metric spaces are quasi-isometric to connected graphs (see [Gro93] p.7 for a closely related fact).

Observation 1.13. Let Y be a geodesic metric space and let $V \subset Y$ be a subset such that for all $y \in Y$ there exists $z \in V$ such that $d(y, z) \leq D$. Let $E \geq 2D + 1$. Let Z be a graph such that
a) the vertex set $\mathcal{V}(Z) = V$
b) the edge set $\mathcal{E}(Z)$ is given by $\{y, z\} \in \mathcal{E}(Z)$ iff $d(y, z) \leq E$.
Define $\psi_Z : Z \to Y$ as follows: $\psi_Z(u) = u$ for $u \in V$. For an edge e of Z choose some $u \in V$ such that e is incident on u and map the interior of e to u under ψ_Z. Then ψ_Z is a quasi-isometry.

A quasi-isometric inverse ψ_Y of ψ_Z can be constructed explicitly as follows. Define $\psi_Y(u) = u$ for $u \in V$. For $y \in Y \setminus V$ choose some $z \in V$ such that $d(y, z) \leq D$ and define $\psi_Y(y) = z$.

A bit more will be necessary in Lemma 1.14 below. Fix some $E_1 > E$ and introduce extra edges in Z for some (not necessarily all) vertices $\{y, z\} \in V$, satisfying $d(y, z) \leq E_1$. The resulting graph, Z_1 say, is quasi-isometric to Z, and hence to Y.

Now, suppose $p : X^{'} \to B^{'}$ is an (f, c, K)-metric bundle. Let d denote the metric on $X^{'}$ and let $d_{B'}$ be the metric on $B^{'}$. Let $V \subset B^{'}$ be a maximal subset such that $u, v \in V, u \neq v$ implies $d_{B'}(u, v) \geq 1$. Construct a graph B as in Observation 1.13 with vertex set V such that $u \neq v \in V$ are connected by an edge iff $d_{B'}(u, v) \leq 3$. As in Observation 1.13, we can define $\psi_B, \psi_{B'}$ to show that $B, B^{'}$ are quasi-isometric.

Next, for all $u \in V$ let $X_u^{'}$ be a maximal subset of the horizontal space F_u such that for $x, y \in X_u^{'}$, $d_u(x, y) \geq 1$. If $u, v \in V$ are connected by an edge in B then note that each point of $X_u^{'}$ is connected to a point of $X_v^{'}$ by a path in $X^{'}$ of length at most $3c+1$. Now construct a graph X with vertex set $\cup_{u\in V} X_u^{'}$ and two vertices $x \neq y$ connected by an edge iff $d(x, y) \leq 6c + 3$ and
a) either $x, y \in X_u^{'}$ for some $u \in V$
b) or $x \in X_u^{'}$ and $y \in X_v^{'}$ with $d_B(u, v) = 1$.

Define $\pi : X \to B$ by sending edges connecting any two vertices of $X_u^{'}$ (for some $u \in V$) to u. Any other edge in X must join vertices $x \in X_u^{'}$ and $y \in X_v^{'}$ for some $X_u^{'}, X_v^{'}$ with $d_B(u, v) = 1$. On any such edge $[x, y]$, p is defined to be an isometry onto the edge $[u, v]$.

Lemma 1.14. *Let X, B, π be as above. Then the map $\pi : X \to B$ gives a metric graph bundle. Further $X, X^{'}$ are quasi-isometric and $B, B^{'}$ are quasi-isometric.*

Proof. By the construction above, π is a simplicial map. Note that $\pi^{-1}(u)$ is a graph with vertex set $X_u^{'}$ and edges of type (a) above. By Observation 1.13, $\pi^{-1}(u)$ is a connected subgraph of X, quasi-isometric to F_u.

Let $x, y \in \pi^{-1}(u)$. To check Condition 1 of the definition of a metric graph bundle, we may assume without loss of generality that $x, y \in X_u^{'}$. If $x, y \in X_u^{'}$ be such that $d_X(x, y) \leq N$, then $d(x, y) \leq N(6c + 3)$. Since $p : X^{'} \to B^{'}$ is an (f, c, K)-metric bundle it follows that $d_u(x, y) \leq f(N(6c + 3))$. Hence $d_{X_u^{'}}(x, y) \leq f(N(6c+3))$, where $d_{X_u^{'}}$ is the metric on $X_u^{'}$ regarded as the vertex set of the graph $\pi^{-1}(u)$. Defining $g(N) = f(N(6c + 3))$, we see that Condition 1 of the definition of a metric graph bundle is satisfied.

Next, if $u, v \in V$ are connected by an edge in B, then for any $x \in X_u^{'}$ there exists $y \in X_v^{'}$ and a path in $X^{'}$ of length at most $3c + 1$ connecting x, y. Since $3c + 1 < 6c + 3$, it follows that there is an edge in X joining x, y. Thus Condition 2(i) of the definition of a metric graph bundle is satisfied. This proves the first assertion of the Lemma.

The above construction gives more. We have already noted that $B, B^{'}$ are quasi-isometric by Observation 1.13. The total spaces $X, X^{'}$ are also quasi-isometric as the following argument shows.

First, $\psi_X : X \to X^{'}$ can be defined as in Observation 1.13 (esp. the last statement) by $\psi_X(x) = x$ for $x \in \cup_{u\in V} X_u^{'}$. Then $p \circ \psi_X = \psi_B \circ \pi$ on $\cup_{u\in V} X_u^{'}$. Then ψ_X is extended over edges by sending any edge to a vertex on which it is incident consistently ensuring that $p \circ \psi_X = \psi_B \circ \pi$.

We need to be a little more careful in defining the quasi-isometric inverse $\psi_{X'}$: $X^{'} \to X$. Suppose $x \in X^{'}$ and let $\psi_{B'}(p(x)) = u \in V$. Choose a point $y \in F_u$ such that x, y can be joined by a path of length at most c (such a path exists by the definition of a metric bundle). Now let $x_1 \in X_u^{'}$ be such that $d_u(x_1, y) \leq 1$. Define $\psi_{X'}(x)$ to be x_1. Note that $d(x, x_1) \leq c + 1$, i.e. $\psi_{X'}$ moves any point through at most a distance $(c+1)$. As in Observation 1.13, $\psi_{X'}$ is a quasi-isometric inverse of ψ_X.

Further, $\pi \circ \psi_{X'}(x) = \pi(x_1) = u = \psi_{B'} \circ p(x)$. Since x is arbitrary, it follows that $\pi \circ \psi_{X'} = \psi_{B'} \circ p$. We note that the maps ψ_X and $\psi_{X'}$ when restricted to horizontal spaces with their respective path metrics give quasi-isometries too (by Observation 1.13). $\square$

Note: In the rest of the paper we shall assume that all the maps $\psi_X, \psi_B, \psi_{X'}, \psi_{B'}$ above are K_1-quasi-isometries. We shall refer to $\pi : X \to B$ above as an *approximating metric graph bundle* of the metric bundle $p : X^{'} \to B^{'}$.

1.2. **Hyperbolic metric spaces.**

We assume that the reader is familiar with the basic definitions and facts about hyperbolic metric spaces [Gro85], [Gd90], [ABC$^+$91]. In this subsection we collect together some of these to fix notions and for later use.

Convention: We shall use **subscripts** for constants to indicate the Lemma/ Proposition/Theorem where they first appear.

If X is a geodesic metric space and $x, y \in X$ then $[x, y]$ will denote a geodesic segment joining x to y. For $x, y, z \in X$ we shall denote by $\triangle xyz$ a geodesic triangle with vertices x, y, z. For $D \geq 0$ and $A \subset X$, $N_D(A) := \{x \in X : d(x, a) \leq D \text{ for some } a \in A\}$ will be called the D-neighborhood of A in X.

Definition 1.15. *Suppose $\Delta x_1x_2x_3 \subset X$ is a geodesic triangle, and let $\delta \geq 0$.*

1. *For all $i \neq j \neq k \neq i$, let $c_k \in [x_i, x_j]$ be such that $d(x_i, c_j) = d(x_i, c_k)$. The points c_i will be called the* **internal points** *of $\Delta x_1x_2x_3$. Note that, for all $i \neq j \neq k \neq i$, $d(x_i, c_j) = \frac{1}{2}\{d(x_i, x_j) + d(x_i, x_k) - d(x_j, x_k)\}$.*
2. *The diameter of the set $\{c_1, c_2, c_3\}$ will be referred to as the* **insize** *of the triangle $\Delta x_1x_2x_3$.*
3. *We say that the triangle $\Delta x_1x_2x_3$ is δ-***slim** *if any side of the triangle is contained in the δ-neighborhood of the union of the other two sides.*
4. *We say that the triangle $\Delta x_1x_2x_3$ is δ-***thin** *if for all $i \neq j \neq k \neq i$ and $p \in [x_i, c_j] \subset [x_i, x_k]$, $q \in [x_i, c_k] \subset [x_i, x_j]$ with $d(p, x_i) = d(q, x_i)$ one has $d(p, q) \leq \delta$.*

Definition 1.16. Gromov inner product: *Let X be any metric space and let $x, y, z \in X$. Then the Gromov inner product of y, z with respect to x, denoted $(y.z)_x$, is defined to be the number $\frac{1}{2}\{d(x, y) + d(x, z) - d(y, z)\}$.*

Definition 1.17. *Let $\delta \geq 0$ and X be a geodesic metric space. We say that X is a δ-hyperbolic metric space if all geodesic triangles in X are δ-slim.*

Lemma 1.18. *(See Proposition 2.1,[ABC$^+$91]) Suppose X is a δ-hyperbolic metric space. Then the following hold:*

1. *All the triangles in X have insize at most 4δ.*
2. *All the triangles in X are 6δ-thin.*

Lemma 1.19. [Gd90] **Stability of quasigeodesics:** *For all $\delta \geq 0$ and $k \geq 1$ there is a constant $D_{1.19} = D_{1.19}(\delta, k)$ such that the following holds:*

Suppose Y is a δ-hyperbolic metric space. Then the Hausdorff distance between a geodesic and a k-quasi-geodesic joining the same pair of end points is less than or equal to $D_{1.19}$.

Definition 1.20. Local quasi-geodesics: *Let X be a metric space and $K \geq 1, \epsilon \geq 0, L > 0$ be constants. A map $f : I \to X$, where $I \subset \mathbb{R}$ is an interval, is said to be a $(K, \epsilon, L)-$ local quasi-geodesic if for all $s, t \in I$ with $|s - t| \leq L$, one has $-\epsilon + (1/K)|s - t| \leq d(f(s), f(t)) \leq \epsilon + K|s - t|$.*

For the following important lemma we refer to Theorem 1.4, Chapter 3,[CDP90]; or Theorem 21, Chapter 5, [Gd90].

Lemma 1.21. Local quasi-geodesic vs global quasi-geodesic: *For all $\delta \geq 0$, $\epsilon \geq 0$ and $K \geq 1$ there are constants $L = L_{1.21}(\delta, K, \epsilon)$, $\lambda = \lambda_{1.21}(\delta, K, \epsilon)$ such that the following holds:*

Suppose X is a δ-hyperbolic metric space. Then any (K, ϵ, L)-local quasi-geodesic in X is a λ-quasi-geodesic.

Lemma 1.22. *For all $\delta \geq 0$, $\epsilon \geq 0$ and $k \geq 1$, there is a constant $D_{1.22} = D_{1.22}(\delta, k, \epsilon)$ such that the following hold:*

(1) Suppose Y is a δ-hyperbolic metric space and $y_1, y_2, y_3 \in Y$. Then, there is a point $y \in Y$ such that $y \in N_{4\delta}([y_i, y_j])$ for any geodesic $[y_i, y_j]$ joining y_i, y_j, $i \neq j \in \{1, 2, 3\}$. We call such a point a barycenter *of the triangle $\triangle y_1 y_2 y_3$.*

(2) Suppose both Y and $Y^{'}$ are δ-hyperbolic metric spaces and $\phi : Y \to Y^{'}$ is a (k, ϵ)-quasi-isometric embedding. If y is a barycenter of $\triangle y_1 y_2 y_3 \subseteq Y$ and $y^{'} \in Y^{'}$ is a barycenter of $\triangle \phi(y_1)\phi(y_2)\phi(y_3) \subseteq Y^{'}$ then $d(y^{'}, \phi(y)) \leq D_{1.22}$, where d is the metric on $Y^{'}$.

Proof. By conclusion (1) of Lemma 1.18 the internal points of $\triangle y_1 y_2 y_3$ are barycenters of $\triangle y_1 y_2 y_3$. This proves part (1) of the lemma.

For (2), first we make the following simple observation: Let $c_i^{'}$ be the internal points of $\triangle \phi(y_1)\phi(y_2)\phi(y_3)$. Suppose $z \in Y^{'}$ is contained in a D-neighborhood of each of the sides of $\triangle \phi(y_1)\phi(y_2)\phi(y_3)$, for some $D \geq 0$. Let $p_i \in [\phi(y_j), \phi(y_k)]$, $i \neq j \neq k \neq i$, be such that $d(p_i, z) \leq D$, $1 \leq i, j, k \leq 3$; then $d(p_i, p_j) \leq 2D$. The computation below shows that $d(c_i^{'}, p_i) \leq 3D$ (we do the computation for $i = 3$ for concreteness).

Set $A_i = \phi(y_i), i = 1, 2, 3$. Then

$$\begin{aligned}
&2d(p_3, c_3^{'})\\
&= 2|d(A_1, c_3^{'}) - d(A_1, p_3)|\\
&= 2|(A_2, A_3)_{A_1} - d(A_1, p_3)|\\
&= |d(A_1, A_3) + d(A_1, A_2) - d(A_2, A_3) - 2d(A_1, p_3)|\\
&= |\{d(A_1, p_2) + d(A_3, p_2)\} + \{d(A_1, p_3) + d(A_2, p_3)\}\\
&\qquad\qquad - \{d(A_3, p_1) + d(A_2, p_1)\} - 2d(A_1, p_3)|\\
&= |\{d(A_1, p_2) - d(A_1, p_3)\} + \{d(A_3, p_2) - d(A_3, p_1)\} + \{d(A_2, p_3) - d(A_2, p_1)\}|\\
&\leq |d(A_1, p_2) - d(A_1, p_3)| + |d(A_3, p_2) - d(A_3, p_1)| + |d(A_2, p_3) - d(A_2, p_1)|\\
&\leq d(p_2, p_3) + d(p_2, p_1) + d(p_3, p_1)\\
&\leq 6D
\end{aligned}$$

Hence $d(p_3, c_3^{'}) \leq 3D$ and thus $d(z, c_i^{'}) \leq 4D$ for all i, $1 \leq i \leq 3$.

Next, since ϕ is a (k, ϵ)-quasi-isometric embedding, it follows that $\phi(y)$ is contained in the $(4k\delta + \epsilon)-$ neighborhood of the image under ϕ of each of the sides $[y_i, y_j]$, $i \neq j$. Also, the image of $[y_i, y_j]$, for all $i \neq j$, is a (k, ϵ)-quasi-geodesic, and hence a $(k + \epsilon)$-quasi-geodesic, joining $\phi(y_i)$, $\phi(y_j)$. By Lemma 1.19, $\phi(y)$ is contained in a $\{(4k\delta+\epsilon)+D_{1.19}(\delta,k+\epsilon)\}-$ neighborhood of each of the sides of $\triangle\phi(y_1)\phi(y_2)\phi(y_3)$. Taking $D_{1.22}(\delta,k,\epsilon):=4.\{(4k\delta+\epsilon)+D_{1.19}(\delta,k+\epsilon)\}$, we are through. $\square$

Definition 1.23. *Let X be a geodesic metric space and let $A\subseteq X$. For $K\geq 0$, we say that A is K-quasiconvex in X if any geodesic with end points in A is contained in the K-neighborhood of A. A subset $A\subset X$ is said to be quasi-convex if it is K-quasi-convex for some K.*

Lemma 1.24. *For each $\delta\geq 0$ and $K\geq 0$ there is a constant $D_{1.24}$=$D_{1.24}(\delta,K)$ such that the following holds:*
Suppose X is a δ-hyperbolic metric space and $V\subseteq U$ are K-quasi-convex subsets of X. Let $x\in X$ and let x_1,x_2 be nearest point projections of x on U and V respectively. Then, for every point $x_1^{'}\in U$, the (arc length parametrization of the) union of the geodesics $[x,x_1]\cup[x_1,x_1^{'}]$ is a $(3+2K)$-quasi-geodesic. Moreover, if x_3 is a nearest point projection of x_1 on V, then $d(x_2,x_3)\leq D_{1.24}$.

Proof. First note that $[x,x_1]\cup[x_1,x_1^{'}]$ is a $(3,2K)$-quasi-geodesic, whence it is a $(3+2K)$-quasi-geodesic.

In particular $[x,x_1]\cup[x_1,x_2]$ is a $(3+2k)-$quasi-geodesic. Hence by Lemma 1.19, there is a point $x_4\in[x,x_2]$ with $d(x_1,x_4)\leq D_{1.19}(\delta,3+2K)=D$, say. Similarly, $[x_1,x_3]\cup[x_3,x_2]$ is a $(3+2K)$-quasi-geodesic and thus there is a point $x_3^{'}\in[x_1,x_2]$ such that $d(x_3,x_3^{'})\leq D$. Using the δ-slimness of $\triangle x_1x_2x_4$, there exists a point $x_3^{''}\in[x_2,x_4]$ such that $d(x_3^{'},x_3^{''})\leq D+\delta$. Thus it follows that $d(x_3,x_3^{''})\leq 2D+\delta$. Since x_2 is a nearest point projection of $x_3^{''}$ on V, we have $d(x_2,x_3^{''})\leq 2D+\delta$. Thus $d(x_2,x_3)\leq d(x_2,x_3^{''})+d(x_3^{''},x_3)\leq 4D+2\delta$. This completes the proof of the lemma. $\square$

Definition 1.25. *Suppose Y is a metric space and $U,V\subset Y$. We say that U,V are* ϵ-separated *if $\inf\{d(y_1,y_2):y_1\in U,y_2\in V\}\geq\epsilon$. A collection of subsets $\{U_\alpha\}$ of Y is said to be* **uniformly separated** *if there exists an $\epsilon>0$ such that any pair of distinct elements of the collection $\{U_\alpha\}$ is ϵ-separated.*

The next lemma is a consequence of the first part of Lemma 1.24.

Lemma 1.26. *Given $\delta\geq 0$ and $K\geq 0$ there are constants $R=R_{1.26}(\delta,K)$ and $D=D_{1.26}(\delta,K)$ such that the following holds:*

Suppose X is a δ-hyperbolic metric space and $U,V\subset X$ are two K-quasiconvex and R-separated subsets. Then there are points $x_0\in U$, $y_0\in V$ such that $[x_0,y_0]\subset N_D([x,y])$, for all $x\in U$ and $y\in V$.

Proof. First consider the set $V_1(\subset V)$ of all nearest point projections from points of U onto V. We claim that V_1 is a set of uniformly bounded diameter for large enough R.

Suppose that $x_1,x_2\in U$ and let $y_1,y_2\in V$ be respectively their nearest point projections. Then by Lemma 1.24, $[x_1,y_1]\cup[y_1,y_2]$ and $[x_2,y_2]\cup[y_2,y_1]$ are K_1-quasi-geodesics for $K_1=(3+2K)$. If $d(y_1,y_2)\geq D_1:=L_{1.21}(\delta,K_1,K_1)$ then the curve $[x_1,y_1]\cup[y_1,y_2]\cup[y_2,x_2]$ is a $\lambda=\lambda_{1.21}(\delta,K_1,K_1)$-quasi-geodesic. Hence every point of this curve is within distance $D_{1.19}(\delta,\lambda)+K$ from a point in U. Choosing $R=D_{1.19}(\delta,\lambda)+K+1$ we are through with the claim.

Choose any point $y_0\in V_1$ and let x_0 be a nearest point projection of y_0 onto U. Let $x\in U$, $y\in V$ be any pair of points and let y_1 be a nearest point projection of

x onto V. Since $y_1 \in V_1$, it follows that $d(y_0, y_1) \leq D_1$. It is now clear that each point of $[x_0, y_0]$ is within a distance $2D_{1.19}(\delta, 3+2K) + D_1 + \delta$ from $[x, y]$. $\square$

The following Lemma [Mit98b] says that quasi-isometries and nearest point projections 'almost commute'.

Lemma 1.27. *(Lemma 3.5 of* [Mit98b]*) For all $\delta \geq 0$ and $k \geq 1$ there is a constant $D_{1.27} = D_{1.27}(\delta, k)$ such that the following holds:*
Suppose $\phi : X \to Y$ is a k-quasi isometric embedding of δ-hyperbolic metric spaces. Let $x, y, z \in X$ and let γ be a geodesic in X joining x, y. Let u be a nearest point projection of z onto γ and suppose v is a nearest point projection of $\phi(z)$ onto a geodesic joining $\phi(x)$ and $\phi(y)$, then $d(v, \phi(u)) \leq D_{1.27}$.

To prove our main theorem, the following characterization of hyperbolicity turns out to be very useful.

Lemma 1.28. *(Proposition 3.5 of* [Ham07]*) Suppose X is a geodesic metric space and there is a collection of rectifiable curves $\{c(x,y) : x, y \in X\}$, one for each pair of distinct points $x, y \in X$, and constants $D_1, D_2 \geq 1$ such that for all $x, y, z \in X$ the following hold:*

1. *If $d(x,y) \leq D_1$ then the length of the curve $c(x,y)$ is less than or equal to D_2.*
2. *If $x^{'}, y^{'} \in c(x,y)$ then the Hausdorff distance between $c(x^{'}, y^{'})$ and the segment of $c(x,y)$ between $x^{'}$ and $y^{'}$ is bounded by D_2.*
3. *The triangle formed by the curves joining any three points in X is D_2-slim: $c(x,y) \subseteq N_{D_2}(c(x,z) \bigcup c(y,z))$.*

Then X is $\delta_{1.28} = \delta_{1.28}(D_1, D_2)$-hyperbolic and each of the curves $c(x,y)$ is a $K_{1.28} = K_{1.28}(D_1, D_2)$-quasi-geodesic in X.

This lemma has the following straightforward corollary, which is a discrete version of Lemma 1.28 (see Observation 1.13 for instance). A *discrete path $c(x,y)$* will refer to a finite sequence of points. The *length* of a discrete path is the sum of the distances between all pairs of successive points in the discrete path. In this context, a triangle will refer to the union of three discrete paths of the form $c(x,y)$, $c(y,z)$, $c(z,x)$.

Corollary 1.29. *Let X be a geodesic metric space and let $X_1 \subset X$ be a discrete set such that $X = N_D(X_1)$ for some $D > 0$. Let $D_1, D_2, D_3, D_4 > 0$ be constants such that the following hold.*
For all $x \neq y \in X_1$, there is a discrete path $c(x,y)$ in X_1 connecting x, y such that:

1. *Distance between successive points of $c(x,y)$ is at most D_1.*
2. *If $d(x,y) \leq D_2$ then the length of the path $c(x,y)$ is at most D_3.*
3. *If $x_1 \neq y_1$ are two points of the discrete path $c(x,y)$, then the Hausdorff distance between the discrete path $c(x_1, y_1)$ and the discrete subpath of $c(x,y)$ connecting x_1, y_1 is at most D_4.*
4. *For any three points $x, y, z \in X_1$, the triangle formed by the paths $c(x,y), c(y,z)$ and $c(x,z)$ is D_4-slim.*

Then X is $\delta = \delta(D, D_1, D_2, D_3, D_4)$-hyperbolic and the discrete paths are $K_{1.29} = K_{1.29}(D, D_1, D_2, D_3, D_4)$-quasi-geodesics in X .

Proof. Given $x, y \in X$ choose $x_1, y_1 \in X_1$ such that $d(x, x_1) \leq D, d(y, y_1) \leq D$. Define a curve $\beta(x, y)$ joining x, y as follows: Let $x_1 = v_1, v_2, \ldots, v_n = y_1$ be the set of successive points on $c(x_1, y_1)$. Join x to x_1, y to y_1 and v_i to v_{i+1}, for $1 \leq i \leq n-1$ by geodesics. The set of curves $\{\beta(x, y)\}$ satisfy the conditions of Lemma 1.28 as $\beta(x, y)$ lies in a $\max\{D, D_1\}-$neighborhood of $c(x, y)$. □

1.3. **Trees of hyperbolic and relatively hyperbolic metric spaces.** We refer to [Far98] for a detailed account of relative hyperbolicity. We also refer to [MR08] for the definitions and results of this subsection.

Suppose (X, d) is a path metric space and let $\mathcal{H} = \{H_\alpha\}$ be a collection of path-connected, uniformly separated subsets of X. Then Farb [Far98] defines the **electric space** (or coned-off space) $\mathcal{E}(X, \mathcal{H})$ corresponding to the pair $(X, \mathcal{H})$ as a metric space which consists of X and a collection of vertices v_α (one for each $H_\alpha \in \mathcal{H}$) such that each point of H_α is joined to (equivalently, coned off at) v_α by an edge of length $\frac{1}{2}$. The sets H_α shall be referred to as *horosphere-like sets* and the vertices v_α as cone-points.

When the collection $\mathcal{H} = \{H_\alpha\}$ is *not necessarily separated*, a slightly modified description is given in [Mj06] and [MR08], where we attach a metric product $H_\alpha \times [0, 1]$ to X, identifying $H_\alpha \times \{0\}$ with $H_\alpha \subset X$ for each $H_\alpha \in \mathcal{H}$, and equip each $H_\alpha \times \{1\}$ with the zero metric. We shall call this construction *electrocution*.

Let $i : X \to \mathcal{E}(X, \mathcal{H})$ denote the natural inclusion of spaces. Then for a path $\gamma \subset X$, the path $i(\gamma)$ lies in $\mathcal{E}(X, \mathcal{H})$. Replacing maximal subsegments $[a, b]$ of $i(\gamma)$ lying in a particular H_α by a path that goes from a to v_α and then from v_α to b, and repeating this for every H_α that $i(\gamma)$ meets we obtain a new path $\hat{\gamma}$. If $\hat{\gamma}$ is an electric geodesic (resp. electric P-quasigeodesic), γ is called a *relative geodesic* (resp. *relative P-quasigeodesic*). We shall usually be concerned with the case that γ is a geodesic/quasigeodesic without backtracking, i.e. paths which do not return to an H_α after leaving it.

Definition 1.30. [Far98] Relative P-quasigeodesics in $(X, \mathcal{H})$ are said to satisfy **bounded region penetration** if, for any two relative P-quasigeodesics without backtracking β, γ, joining $x, y \in X$, there exists $B = B(P) \geq 0$ such that
Similar Intersection Patterns 1: if precisely one of $\{\beta, \gamma\}$ meets a horosphere-like set H_α, then the length (measured in the intrinsic path-metric on H_α) from the first (entry) point to the last (exit) point (of the relevant path) is at most B.
Similar Intersection Patterns 2: if both $\{\beta, \gamma\}$ meet some H_α then the length (measured in the intrinsic path-metric on H_α) from the entry point of β to that of γ is at most B; similarly for exit points.

X is *strongly hyperbolic relative to the collection* $\mathcal{H}$ if
a) $\mathcal{E}(X, \mathcal{H})$ is hyperbolic, and
b) Relative quasigeodesics satisfy the bounded region penetration property.

The next notion is based on Bestvina-Feighn's seminal work [BF92]. The notions we use here are the adaptations used in [MR08].

Definition 1.31. *$P : X \to T$ is said to be a tree of geodesic metric spaces satisfying the q(uasi) i(sometrically) embedded condition if the geodesic metric space (X, d) admits a map $P : X \to T$ onto a simplicial tree T, such that there exist $\epsilon \geq 0$ and $K \geq 1$ satisfying the following:*
1) For all vertices $v \in T$, $X_v = P^{-1}(v) \subset X$ with the induced path metric d_{X_v} is a geodesic metric space. Further, the inclusions $i_v : X_v \to X$ are uniformly

proper, i.e. for all $M > 0$*, there exists* $N > 0$ *such that* $d(i_v(x), i_v(y)) \leq M$ *implies* $d_{X_v}(x, y) \leq N$ *for all* $v \in T$ *and* $x, y \in X_v$*.*
2) Let e *be an edge of* T *with initial and final vertices* v_1 *and* v_2 *respectively. Let* X_e *be the pre-image under* P *of the mid-point of* e*. There exist continuous maps* $f_e : X_e \times [0, 1] \to X$*, such that* $f_e|_{X_e \times (0,1)}$ *is an isometry onto the pre-image of the interior of* e *equipped with the path metric. Further,* f_e *is fiber-preserving, i.e. projection to the second co-ordinate in* $X_e \times [0, 1]$ *corresponds via* f_e *to projection to the tree* $P : X \to T$*.*
3) $f_e|_{X_e \times \{0\}}$ *and* $f_e|_{X_e \times \{1\}}$ *are* (K, ϵ)*-quasi-isometric embeddings into* X_{v_1} *and* X_{v_2} *respectively.* $f_e|_{X_e \times \{0\}}$ *and* $f_e|_{X_e \times \{1\}}$ *will occasionally be referred to as* f_{e,v_1} *and* f_{e,v_2} *respectively.*

X_v, X_e are referred to as vertex and edge spaces respectively. A tree of spaces as in Definition 1.31 above is said to be a *tree of hyperbolic metric spaces*, if there exists $\delta > 0$ such that X_v, X_e are all δ-hyperbolic for all vertices v and edges e of T.

Definition 1.32. *A tree* $P : X \to T$ *of geodesic metric spaces is said to be a tree of relatively hyperbolic metric spaces if in addition to the conditions of Definition 1.31, we have the following:*
4) Each vertex (or edge) space X_v *(or* X_e*) is strongly hyperbolic relative to a collection* $\mathcal{H}_v$ *(or* $\mathcal{H}_e$*)*
5) the maps f_{e,v_i} *above (*$i = 1, 2$*) are* **strictly type-preserving***, i.e.* $f^{-1}_{e,v_i}(H_{v_i,\alpha})$*,* $i = 1, 2$ *(for any* $H_{v_i,\alpha} \in \mathcal{H}_{v_i}$*) is either empty or some* $H_{e,\beta} \in \mathcal{H}_e$*. Also, for all* $H_{e,\beta} \in \mathcal{H}_e$*, and any end-point* v *of* e*, there exists* $H_{v,\alpha}$*, such that* $f_{e,v}(H_{e,\beta}) \subset H_{v,\alpha}$*.*
6) There exists $\delta > 0$ *such that each* $\mathcal{E}(X_v, \mathcal{H}_v)$ *is* δ*-hyperbolic.*
7) The induced maps (see below) of the coned-off edge spaces into the coned-off vertex spaces $\widehat{f_{e,v_i}} : \mathcal{E}(X_e, \mathcal{H}_e) \to \mathcal{E}(X_{v_i}, \mathcal{H}_{v_i})$ *(*$i = 1, 2$*) are uniform quasi-isometries. This is called the* **qi-preserving electrocution condition***.*

d_v and d_e will denote path metrics on X_v and X_e respectively. i_v, i_e will denote inclusion of X_v, X_e respectively into X.

For a tree of relatively hyperbolic spaces, the sets $H_{v\alpha}$ and $H_{e\alpha}$ will be referred to as **horosphere-like vertex sets and edge sets** respectively.

Given the tree of spaces with vertex spaces X_v and edge spaces X_e, there exists a naturally associated tree with vertex spaces $\mathcal{E}(X_v, \mathcal{H}_v)$ and edge spaces $\mathcal{E}(X_e, \mathcal{H}_e)$, obtained by simply coning off the respective horosphere like sets. Condition (5) of the above definition ensures that we have natural inclusion maps of edge spaces $\mathcal{E}(X_e, \mathcal{H}_e)$ into adjacent vertex spaces $\mathcal{E}(X_v, \mathcal{H}_v)$. These are the maps referred to as *induced maps* in Condition (7) above.

The resulting tree of coned-off spaces will be called the **induced tree of coned-off spaces** and will be denoted as $\widehat{X}$.

Definition 1.33. *The* **cone locus** *of the induced tree (*T*) of coned-off spaces,* $\widehat{X}$*, is the graph whose vertex set* $\mathcal{V}$ *consists of horosphere like vertex sets and edge set* $\mathcal{E}$ *consists of horosphere like edge sets such that an edge* $H_{e,\beta} \in \mathcal{H}_e \subset \mathcal{E}$ *is incident on a vertex* $H_{v_i,\alpha} \in \mathcal{H}_{v_i} \subset \mathcal{V}$ *iff* $f_{e,v}(H_{e,\beta}) \subset H_{v,\alpha}$*.*
A connected component of the cone-locus will be called a **maximal cone-subtree***. The collection of maximal cone-subtrees will be denoted by* $\mathcal{T}$ *and elements of* $\mathcal{T}$ *will be denoted as* T_α*.*
For each maximal cone-subtree T_α*, we define the associated* **maximal cone-subtree**

of horosphere-like spaces C_α *to be the tree of metric spaces whose vertex and edge spaces are the horosphere like vertex and edge sets* $H_{v\alpha}$, $H_{e\alpha}$, $v \in \mathcal{V}(T_\alpha)$, $e \in \mathcal{E}(T_\alpha)$, *along with the restrictions of the maps* f_e *to* $H_{e\alpha} \times \{0,1\}$. *The collection of* C_α*'s will be denoted as* $\mathcal{C}$.

The next definition is based on [BF92] again.

Definition 1.34. *A disk* $f : [-m,m]\times I \to X$ *is a* **hallway** *of length* $2m$ *if it satisfies:*
1) $f^{-1}(\cup X_e : e \in \mathrm{Edge}(T)) = \{-m, \cdots, m\}\times I$, *where* $\mathrm{Edge}(T)$ *denotes the collection of mid-points of the edge-set of* T.
2) f *maps* $i\times I$ *to a geodesic in* X_e *for some edge space.*
3) f *is transverse, relative to condition (1), to* $\cup_e X_e$, *i.e. for all* $i \in \{-m, \cdots, m\}$ *and* $t \in I$, *if* $f|_{N(i,\frac{1}{4})\times\{t\}}$ *is an isometric embedding. Here* $N(i, \frac{1}{4})$ *denotes the* $\frac{1}{4}$ *neighborhood of* i *in* $[-m,m]$.

Condition (3) above is the adaptation in our context of Condition (2) of [BF92] p.87 and simply says that a hallway transversely cuts across the collection of edge spaces.

Definition 1.35. *A hallway is* ρ**-thin** *if* $d(f(i,t), f(i+1,t)) \le \rho$ *for all* i, t.
A hallway is λ**-hyperbolic** *if*

$$\lambda l(f(\{0\} \times I)) \le \ max\{l(f(\{-m\} \times I)), l(f(\{m\} \times I))\}$$

where $l(\sigma)$ *denotes the length of the path* σ.
The **girth** *of the hallway is the quantity* $l(f(\{0\} \times I))$.
A hallway is **essential** *if the edge path in* T *resulting from projecting* $f : [-m,m]\times I(\subset X)$ *onto* T *does not backtrack, and is therefore a geodesic segment in the tree* T.
Hallways flare condition: *The tree of spaces,* X, *is said to satisfy the hallways flare condition if there are numbers* $\lambda > 1$ *and* $m \ge 1$ *such that for all* ρ *there is a constant* $H(\rho)$ *such that any* ρ*-thin essential hallway of length* $2m$ *and girth at least* H *is* λ*-hyperbolic.*

Definition 1.36. *An essential hallway of length* $2m$ *is* **cone-bounded** *if*
a) $f(i \times \partial I) = f(i \times \{0,1\})$ *lies in the cone-locus for* $i = \{-m, \cdots, m\}$.
b) $f(i \times \{0\})$ *and* $f(i \times \{1\})$ *lie in different components of the cone-locus.*
The tree of spaces, X, *is said to satisfy the* **cone-bounded hallways strictly flaring condition** *if for all* $\rho > 0$, *there exists* $\lambda > 1$ *and* $m \ge 1$ *such that any cone-bounded* $\rho-$*thin essential hallway of length* $2m$ *is* λ*-hyperbolic.*

Note that the last condition requires all cone-bounded $\rho-$thin essential hallways to flare (not just those of girth less than H as in Definition 1.35). The following theorem is one of the main results of [MR08].

Theorem 1.37. [MR08] *Let* X *be a tree* (T) *of strongly relatively hyperbolic spaces satisfying*

1. *the qi-embedded condition.*
2. *the strictly type-preserving condition.*
3. *the qi-preserving electrocution condition.*
4. *the induced tree of coned-off spaces satisfies the hallways flare condition.*
5. *the cone-bounded hallways strictly flaring condition.*

Then X is (strongly) hyperbolic relative to the family $\mathcal{C}$ of maximal cone-subtrees of horosphere-like spaces.

Note: In [MR08] the definition of cone-bounded hallways does not include Condition (b) of Definition 1.36. However the proof in [MR08] (cf. Proposition 4.4 of [MR08]) gives Theorem 1.37.

Definition 1.38. Partial Electrocution: *Let $(X, \mathcal{H}, \mathcal{G}, \mathcal{L})$ be an ordered quadruple and let $\delta, \epsilon \geq 0, K \geq 1$ be constants such that the following hold:*

1. *X is (strongly) hyperbolic relative to a collection of subsets H_α.*
2. *For each H_α there is a δ-hyperbolic metric space L_α and a map $g_\alpha : H_\alpha \to L_\alpha$ such that $d_{L_\alpha}(g_\alpha(x), g_\alpha(y)) \leq K d_{H_\alpha}(x, y) + \epsilon$ for all $x, y \in H_\alpha$. We denote the collection of such g_α's as $\mathcal{G}$.*

The **partially electrocuted space** *or* partially coned off space *corresponding to $(X, \mathcal{H}, \mathcal{G}, \mathcal{L})$ is the quotient metric space $(\hat{X}, d_{pel})$ obtained from X by attaching the metric mapping cylinders for the maps $g_\alpha : H_\alpha \to L_\alpha$, where d_{pel} denotes the resulting partially electrocuted metric. (The metric mapping cylinder for a map $g : A \to B$ is the quotient metric space obtained as a quotient space of the disjoint union of the metric product $A \times [0, 1]$ and B, by identifying $(a, 1) \in A \times \{1\}$ with $g(a) \in B$.)*

Lemma 1.39. [MR08] *(see also Lemmas 1.20. 1.21 of* [MP11]*) $(\hat{X}, d_{pel})$ is a hyperbolic metric space and the sets L_α are uniformly quasiconvex in $\hat{X}$.*

We end this subsection with a proposition, a special case of which is due to Hamenstadt [Ham05], where the tree is taken to be $\mathbb{R}$ with vertex set $\mathbb{Z}$. We give a different proof below as our proof applies in a more general context. We need the following definition to state the proposition.

Definition 1.40. *Suppose Y is a δ-hyperbolic metric space and U_1, U_2 are two quasi-convex subsets. Let $D > 0$.*
We say that U_1, U_2 are mutually D-cobounded, or simply D-cobounded, if any nearest point projection of U_1 onto U_2 has diameter at most D and vice versa.

The essential feature of D-coboundedness that we shall use below is the fact that if $a \in U_1, b \in U_2$ such that $d(a, b) = d(U_1, U_2)$, and $[c, a] \subset U_1$, $[b, d] \subset U_2$ are K-quasigeodesics, then $[c, a] \cup [a, b] \cup [b, d]$ is a $K_1 (= K_1(K, \delta))$-quasigeodesic (see Lemma 3.3 of [Mit98b] for instance).

Proposition 1.41. *Given $K \geq 1$ and $\delta, \epsilon, D > 0$, there exist $\delta^{'}, k^{'} \geq 0$ such that the following holds.*
Suppose Y is a tree of δ-hyperbolic metric spaces satisfying the K-qi embedded condition such that the images of the edge spaces in the vertex spaces are mutually D-cobounded. Then Y is a $\delta^{'}$-hyperbolic metric space and all the vertex spaces and edge spaces are $k^{'}$-quasiconvex in Y.

Proof. First of all we note that by [Bow97] (Section 7, esp. Proposition 7.4, Lemma 7.5, Proposition 7.12; see also Lemma 3.4 of [Mj11]) the vertex spaces are strongly hyperbolic relative to the images of edge spaces. Hence Y can be thought of as a tree of relatively hyperbolic metric spaces whose horosphere like edge sets and vertex sets are respectively the whole of the edge spaces and the images of the edge spaces in the vertex spaces respectively. Hence conditions (1)-(2) of Theorem 1.37 are satisfied in this case.

Edge spaces after electrocution become points. Vertex spaces after electrocution become hyperbolic metric spaces. Inclusion of points into spaces being trivially qi-embeddings, condition (3) of Theorem 1.37 is satisfied.

Next any essential hallway of length greater than two in $\hat{Y}$, the induced tree of coned off spaces, must have girth at most one. This is because all edge spaces have diameter one after electrocution. Hence Condition (4) of Theorem 1.37 is trivially satisfied by choosing the threshold value H of the girth to be 2.

Finally, since cone-bounded hallways must have length two or more, it follows that in the present situation cone-bounded hallways do not exist due to Condition (b) in Definition 1.36 and the fact that entire edge spaces are part of the cone locus. Hence Condition (5) of Theorem 1.37 is vacuously satisfied.

Finally, the family $\mathcal{C}$ of maximal cone-subtrees of horosphere-like spaces are precisely the edge spaces.

Hence Y is strongly hyperbolic relative to the edge spaces.

The edge spaces are uniformly hyperbolic with respect to the induced length metric from Y. Hence, by Lemma 1.39, we see that when the maps g_α are taken to be identity maps of the edge spaces, the partially electrocuted space is hyperbolic. This space is clearly quasi-isometric to Y. Hence the result. □

As an application of this proposition we have the following corollary which can be thought of as a 'discrete' or 'graph' version of Proposition 1.41.

Corollary 1.42. *Given $\delta, D, D_1, K \geq 1$, there exists $D_{1.42}$ such that the following holds.*
Suppose X is a connected graph and X_i, $0 \leq i \leq n$, are connected subgraphs such that the following conditions hold.
(1) *All the spaces X_i are δ-hyperbolic with respect to the induced path metric from X.*
(2) *$X_i \cap X_j \neq \emptyset$ iff $|i-j| \leq 1$.*
(3) *For all i, $X_i \cap X_{i+1}$ contains a connected subgraph Y_i and is contained in the D-neighborhood of Y_i in X_i as well as X_{i+1}.*
(4) *The inclusions $Y_i \hookrightarrow X_i$, $Y_i \hookrightarrow X_{i+1}$ are K-quasi-isometric embeddings.*
(5) *Y_i and Y_{i+1} are D_1-cobounded in X_{i+1}.*
Then the space X is $D_{1.42}(= D_{1.42}(\delta, D, D_1, K))$-hyperbolic.

Proof. First construct a new graph $X^{'}$ with the same vertex set as X and edge-set $\mathcal{E}(X^{'}) = \{\{u,v\} : u \neq v; d_X(u,v) \leq D\}$. Note that X is a subgraph of $X^{'}$. Let us denote by $X_i^{'}$ the new graph obtained from X_i after the introduction of the new edges. Let $Y_i^{'} := X_i^{'} \cap X_{i+1}^{'}$. Note that $Y_i^{'}$ is a connected graph by Condition (3).

Now X, X_i, Y_i are quasi-isometric to $X^{'}, X_i^{'}$ and $Y_i^{'}$ respectively (Observation 1.13). Then there exist $\delta^{'}, K^{'}, D_1^{'}$ such that $X_i^{'}$ is $\delta^{'}$ hyperbolic; the inclusion maps $Y_i^{'} \hookrightarrow X_i^{'}$, $Y_i^{'} \hookrightarrow X_{i+1}^{'}$ are $K^{'}$-qi embeddings; and $Y_i^{'}$ and $Y_{i+1}^{'}$ are $D_1^{'}$-cobounded in $X_{i+1}^{'}$ for all i. Now we construct a tree of metric spaces where the underlying tree is the interval $[0,n]$ with vertices the integer points.

Let $e = [i, i+1]$. For each $Y_i^{'}$ construct $Y_i^{'} \times [0,1]$. Now identify $x \times \{0\}$ with $x \in X_i^{'}$ and $x \times \{1\}$ with $x \in X_{i+1}^{'}$ for all $x \in \mathcal{V}(Y_i^{'})$. Extend the identification linearly over edges of $Y_i^{'}$. Performing this identification for all i, we see that $X^{'}$ (and hence X) is quasi-isometric to a tree of metric spaces satisfying all the conditions of Proposition 1.41. □

2. QI Sections

2.1. **Existence of qi sections.** The main result (Proposition 2.10) of this subsection is that qi sections exist for a large class of examples of metric graph bundles $p : X \to B$. This is *the* crucial ingredient in the proof of our main theorem 4.10. The basic idea of the proof of Proposition 2.10 runs as follows:
We assume that the horizontal spaces F_b, $b \in B$ in our metric graph bundle are uniformly hyperbolic and the barycenter maps $\phi_b : \partial^3 F_b \to F_b$, sending a triple of distinct points on the boundary ∂F_b to the barycenter of an ideal triangle with the three points as vertices, are uniformly coarsely surjective, i.e. F_b lies in a uniformly bounded neighborhood of the image of ϕ_b. For simplicity, suppose we have $x \in F_v$, $v \in B$ and there is a triple $\xi = (\xi_1, \xi_2, \xi_3)$ such that $\phi_v(\xi) = x$. Fix one such triple. 'Flow' this triple to the boundaries of all other horizontal spaces F_w by maps induced by quasi-isometries $f_{vw} : F_v \to F_w$. These maps are coarsely unique and are naturally associated to any given metric graph bundle. Consider the barycenters of the ideal triangles formed by the flowed triples $(\partial(f_{vw})\xi_1, \partial(f_{vw})\xi_2, \partial(f_{vw}\xi_3)$. The collection of all these barycenters (as w ranges over B) is then a section through x. The proof that this is indeed a qi section hinges on the fact that for any three points $u, v, w \in B$, the quasi-isometries f_{uv} and $f_{wv} \circ f_{uw}$ are at a bounded distance *(depending on u, v, w)* from each other, and hence the induced boundary maps satisfy the equality $\partial(f_{uv}) = \partial(f_{wv}) \circ \partial(f_{uw})$.

As an application of the proof of this result we recover an important lemma due to Mosher (see Theorem 2.11). It should be noted that though a basic ingredient for both Mosher's proof and ours is the notion of a 'barycenter', we do not have a group action on the boundaries of fiber spaces in our context. Mosher extracts his qi-section from an action of the whole group on the boundary of the normal subgroup.

Definition 2.1. Sequential Boundary*(See Chapter* 4,[ABC+91]*) Let X be a δ-hyperbolic metric space. A sequence of points $\{x_n\}$ in X is said to converge to infinity, written $x_n \to \infty$, if for some (and hence any) point $p \in X$, $lim_{m,n\to\infty}(x_m.x_n)_p = \infty$.*

Define an equivalence relation on the set of all sequences in X converging to infinity, by setting $\{x_n\} \sim \{y_n\}$ iff $lim_{n\to\infty}(x_n.y_n)_p = \infty$. The set of all equivalence classes $\{[\{x_n\}] : x_n \to \infty\}$ will be denoted by ∂X and will be referred to as the sequential boundary of X or simply the boundary of X .

Suppose $\{x_n\}$ is a sequence of points in X and $x_n \to \infty$. We shall write $x_n \to \xi \in \partial X$ to mean that $\xi = [\{x_n\}]$. The boundary ∂X comes equipped with a natural 'visual' topology [Gd90].

Suppose $f : X \to Y$ is a (k, ϵ)-quasi-isometric embedding of hyperbolic metric spaces and $\xi = [\{x_n\}] \in \partial X$. Then $f(x_n) \to \infty$. Setting $\partial(f)(\xi) := [\{f(x_n)\}]$ gives a well defined map $\partial(f) : \partial X \to \partial Y$. The next lemma collects together standard properties of such maps.

Lemma 2.2. *Suppose $f : X \to Y$ is a (k, ϵ)-quasi-isometric embedding of δ-hyperbolic metric spaces. Then we have an induced map $\partial X \to \partial Y$, denoted by $\partial(f)$, such that the following hold.*

(1) *If $I_X : X \to X$ is the identity map then $\partial(I_X)$ is the identity map on the sequential boundary of X.*

(2) *If* $f : X \to Y$ *and* $g : Y \to Z$ *are two* (k, ϵ)*-quasi-isometric embeddings then* $\partial(g \circ f) = \partial(g) \circ \partial(f)$

(3) *If* $f, g : X \to Y$ *are two* (k, ϵ)*-quasi-isometric embeddings such that one has* $sup_{x \in X} d(f(x), g(x)) < \infty$ *then* $\partial(f) = \partial(g)$*.*

(4) *If* $f : X \to Y$ *is a quasi-isometry then* $\partial(f) : \partial X \to \partial Y$ *is a homeomorphism.*

The next lemma is a straightforward consequence of the stability of quasi-geodesics (Lemma 1.19) in hyperbolic metric spaces.

Lemma 2.3. *Let* X *be a* δ*-hyperbolic metric space and let* $\gamma : [0, \infty) \to X$ *be a* (K, ϵ)*-quasi geodesic ray. Let* $\{t_n\}$ *be any sequence of non-negative real numbers tending to* ∞*; then* $\gamma(t_n) \to \infty$ *and the point of* ∂X *represented by* $\{\gamma(t_n)\}$ *is independent of the sequence* $\{t_n\}$*.*

The point of ∂X determined by a quasi-geodesic ray γ will be denoted by $\gamma(\infty)$. The next lemma constructs quasigeodesic rays joining points in X to points in ∂X as well as bi-infinite quasigeodesics joining pairs of points in ∂X. While this is standard for proper X [Gd90], ready references for arbitrary (non-proper) X are a bit difficult to come by, and we include a proof for completeness.

Lemma 2.4. *For any* $\delta \geq 0$ *there is a constant* $K = K_{2.4}(\delta)$ *such that the following holds:*

Suppose X *is a* δ*-hyperbolic metric space.*

(1) *Given any point* $\xi \in \partial X$ *and* $p \in X$ *there is a* K*-quasi-geodesic ray* $\gamma : [0, \infty) \to X$ *of* X *with* $\gamma(0) = p$ *and* $\gamma(\infty) = \xi$*.*

(2) *Given two points* $\xi_1 \neq \xi_2 \in \partial X$ *there is a* K*-quasi-geodesic line* $\alpha : \mathbb{R} \to X$ *with* $\alpha(-\infty) = \xi_1$ *and* $\alpha(\infty) = \xi_2$*.*

Terminology: Any quasi-geodesic ray as in (1) of the above lemma will be referred to as a quasi-geodesic ray joining the points p and ξ. Similarly any quasi-geodesic as in (2) of the above lemma will be referred to as a quasi-geodesic line joining the points ξ_1 and ξ_2.

Proof of Lemma 2.4: (1). We shall inductively construct a suitable sequence of points $\{p_n\}$ such that $p_n \to \xi$ and finally show that the union $\cup[p_n, p_{n+1}]$, of the geodesic segments $[p_n, p_{n+1}]$, is a uniform quasi-geodesic. Suppose $x_n \to \xi$, $x_n \in X$, for all n. Fix $N \geq 1$ and let $p_0 = p$. Since $x_n \to \infty$ we can find a positive integer $n_1 \in \mathbb{N}$ such that $(x_i.x_j)_{p_0} \geq N$ for all $i, j \geq n_1$. Let $[p_0, x_{n_1}]$ be a geodesic joining p_0 and x_{n_1}. Choose $p_1 \in [p_0, x_{n_1}]$ such that $d(p_0, p_1) = N$. Now suppose p_l has been constructed. To construct p_{l+1}, let $n_{l+1} \geq max\{n_k : 1 \leq k \leq l\}$ be an integer such that $(x_i.x_j)_{p_l} \geq (l+1)N$ for all $i, j \geq n_{l+1}$. Choose $p_{l+1} \in [p_l, x_{n_{l+1}}]$ such that $d(p_l, p_{l+1}) = (l+1)N$. Now, let α_N be the arc length parametrization of the concatenation of the geodesic segments $[p_i, p_{i+1}]$, $i \in \mathbb{Z}^+$.

Claim: For $N > max\{7\delta + 1, \frac{1}{3}L_{1.21}(\delta, 1, 42\delta)\}$, α_N is a $\lambda_{1.21}(\delta, 1, 42\delta)$-quasi-geodesic.



First we show that $[p_i, p_{i+1}] \cup [p_{i+1}, p_{i+2}]$ is a uniform quasi-geodesic for each i. Let $n > n_{i+2}$. Join p_i with x_n. Since $(x_n.x_{n_{i+1}})_{p_i} \geq (i+1)N$ and triangles in X are 6δ-thin by Lemma 1.18(2), we can find a point $q_i \in [p_i, x_n]$ such that $d(p_i, q_i) = (i+1)N$ and $d(p_{i+1}, q_i) \leq 6\delta$. Similarly there is a point $q_{i+1} \in [p_{i+1}, x_n]$ such that $d(p_{i+1}, q_{i+1}) = (i+2)N$ and $d(p_{i+2}, q_{i+1}) \leq 6\delta$.

Consider the triangle $\Delta p_i p_{i+1} x_n$. The point $q_{i+1} \in [p_{i+1}, x_n]$ is contained in a δ-neighborhood of $[p_i, p_{i+1}] \cup [p_i, x_n]$. Hence there exists $r_i \in [p_i, x_n] \cup [p_i, p_{i+1}]$ such that $d(r_i, q_{i+1}) \leq \delta$. Since $d(q_{i+1}, p_{i+1}) = (i+2)N$, it follows from the triangle inequality that $d(r_i, q_i) \geq d(q_{i+1}, p_{i+1}) - d(r_i, q_{i+1}) - d(p_{i+1}, q_i) \geq (i+2)N - \delta - 6\delta$. Again, since $N > 7\delta + 1$, it follows that $d(r_i, q_i) \geq (i+1)N + 1$ and hence $r_i \notin [p_i, q_i] (\subset [p_i, x_n])$.

Next, we note that $r_i \notin [p_i, p_{i+1}]$. Else suppose $r_i \in [p_i, p_{i+1}]$. Then $(i+1)N = d(p_i, p_{i+1}) \geq d(r_i, p_{i+1}) \geq d(p_{i+1}, q_{i+1}) - d(r_i, q_{i+1}) = (i+2)N - 6\delta$. This is a contradiction since $N > 7\delta + 1$. Thus $r_i \in [q_i, x_n] \subset [p_i, x_n]$. Also note that $d(p_{i+2}, r_i) \leq d(p_{i+2}, q_{i+1}) + d(q_{i+1}, r_i) \leq 7\delta$.

We now show that $[p_i, p_{i+1}] \cup [p_{i+1}, p_{i+2}]$ is a $(1, 42\delta)$-quasi-geodesic of length at least $3N$ for $N > 7\delta + 1$. Suppose $x \in [p_i, p_{i+1}]$ and $y \in [p_{i+1}, p_{i+2}]$. It is enough to show that $|d(x, p_{i+1}) + d(p_{i+1}, y) - d(x, y)| \leq 42\delta$.

For the δ-slim triangle $\triangle p_i q_i p_{i+1}$, there exists $u \in [p_i, q_i]$ such that $d(x, u) \leq 7\delta$. Similarly for $\triangle p_{i+1} q_i r_i$ and $\triangle p_{i+1} p_{i+2} r_i$, there exists $v \in [q_i, r_i]$ such that $d(y, v) \leq 8\delta$ (the precise constant is obtained by a routine computation).

We have the following inequalities:
$|d(x, p_{i+1}) - d(u, q_i)| \leq d(x, u) + d(p_{i+1}, q_i) \leq 6\delta + 7\delta = 13\delta$,
$|d(p_{i+1}, y) - d(q_i, v)| \leq d(p_{i+1}, q_i) + d(y, v) \leq 7\delta + 8\delta = 15\delta$,
and $|d(u, v) - d(x, y)| \leq d(x, u) + d(y, v) \leq 6\delta + 8\delta = 14\delta$.
Hence $|d(x, p_{i+1}) + d(p_{i+1}, y) - d(x, y)| \leq |\{d(x, p_{i+1}) - d(u, q_i)\} + \{d(p_{i+1}, y) - d(q_i, v)\} + \{d(u, v) - d(x, y)\}| \leq 42\delta$ and we are done.

The claim follows from Lemma 1.21.

Next we show that $\gamma(\infty) = \xi$. For this, by Lemma 2.3, we just need to check that $\{p_n\} \sim \{x_n\}$. Again, to show this, it is enough to check that $\{p_k\} \sim \{x_{n_{k-1}}\}$, i.e. $lim_{k\to\infty}(p_k.x_{n_{k-1}})_p = \infty$. By the above proof we know that $(\cup_{i=1}^{k-1}[p_{i-1}, p_i]) \bigcup [p_{k-1}, x_{n_{k-1}}]$ is a uniform quasi-geodesic. Thus, by stability of quasi-geodesics (Lemma 1.19), we can find a constant D depending only on δ such that there is a point $u \in [p, x_{n_{k-1}}]$ with $d(p_{k-1}, u) \leq D$; similarly there is a point $v \in [p, p_k]$ such that $d(p_{k-1}, v) \leq D$. Therefore, we have $d(u, v) \leq 2D$ and $(p_k.x_{n_{k-1}})_p \geq (u.v)_p \geq d(p, u) - d(u, v) \geq d(p, p_{k-1}) - d(u, p_{k-1}) - d(u, v) \geq d(p, p_{k-1}) - 3D$. As $lim_{k\to\infty} d(p, p_k) = \infty$, we have $lim_{k\to\infty}(p_k.x_{n_{k-1}})_p = \infty$. Therefore, the proof is complete by taking $K_{2.4}(\delta) \geq \lambda_{1.21}(\delta, 1, 42\delta)$.

(2) Let $\lambda := \lambda_{1.21}(\delta, 1, 42\delta)$, and $D_1 := D_{1.19}(\delta, \lambda)$. Now, using the proof of (1), we can construct two λ-quasi-geodesic rays γ_1, γ_2, parametrized by arc length, joining a point $p \in X$ to ξ_1 and ξ_2 respectively. Clearly, $sup\{(x.y)_p : x \in \gamma_1,\ y \in \gamma_2\} < \infty$, else there exist $x_n \in \gamma_1$, $y_n \in \gamma_2$, $n \in \mathbb{N}$, such that $(x_n.y_n)_p \to \infty$. Since $x_n \to \gamma_1(\infty) = \xi_1$ and $y_n \to \gamma_2(\infty) = \xi_2$ by Lemma 2.3, this contradicts the fact that $\xi_1 \neq \xi_2$.

Let $N_1 = sup\{(x.y)_p : x \in \gamma_1, y \in \gamma_2\}$. Let $x_i \in \gamma_i$, $i = 1, 2$, be such that $(x_1.x_2)_p \geq N_1 - 1$. Let $u_i \in [p, x_i]$, $i = 1, 2$ be internal points of $\Delta p x_1 x_2$. By Lemma 1.19 we can find $p_i \in \widehat{px_i}$ such that $d(p_i, u_i) \leq D_1$, $i = 1, 2$. Now, let $\gamma_i^{'} \subset \gamma_i$ be the quasi-geodesic subray starting from p_i, for $i = 1, 2$. We intend to show that the arc length parametrization of the concatenation of $\gamma_1^{'}$, $\gamma_2^{'}$ and a geodesic segment $[p_1, p_2]$ joining p_1, p_2 is a uniform quasi-geodesic (see figure below).



Suppose $y_i \in \gamma_i^{'}$, $i = 1, 2$ and $y_3 \in [p_1, p_2]$. It suffices to find $K \geq 1$ and $\epsilon \geq 0$ independent of y_1, y_2, y_3 such that the following conditions are satisfied.
Condition (1) $l(\widehat{p_1y_1}) + l(\widehat{p_2y_2}) + d(p_1, p_2) \leq Kd(y_1, y_2) + \epsilon$,
Condition (2) $l(\widehat{p_1y_1}) + d(p_1, y_3) \leq Kd(y_1, y_3) + \epsilon$, and
Condition (3) $l(\widehat{p_2y_2}) + d(y_2, y_3) \leq Kd(y_2, y_3) + \epsilon$,
where $\widehat{p_iy_i}$ is the subsegement of γ_i between p_i and y_i for $i = 1, 2$; also for a rectifiable curve segment α, $l(\alpha)$ denotes the length of the curve α. Since the proofs of Conditions (2) and (3) are similar we shall give proofs of Conditions (1) and (2) below.

First of all, we note that $d(u_1, u_2) \leq 4\delta$ by Lemma 1.18 and hence $d(p_1, p_2) \leq d(p_1, u_1) + d(p_2, u_2) + d(u_1, u_2) \leq 2D_1 + 4\delta = D_2$, say. By Lemma 1.19 we can find $z_i \in [p, y_i]$ such that $d(p_i, z_i) \leq D_1$, $i = 1, 2$.

We shall first show that the difference between $(y_1.y_2)_p$ and $(p_1.p_2)_p$ is small. Note that $(y_1.y_2)_p \geq (z_1.z_2)_p \geq (p_1.p_2)_p - \{d(p_1, z_1) + d(p_2, z_2)\} \geq (p_1.p_2)_p - 2D_1$. Also, $|(x_1.x_2)_p - (u_1.u_2)_p| = |d(p, u_1) - (u_1.u_2)_p| = d(u_1, u_2)/2 \leq 2\delta$ and $|(p_1.p_2)_p - (u_1.u_2)_p| \leq d(p_1, u_1) + d(p_2, u_2) \leq 2D_1$. Thus $|(x_1.x_2)_p - (p_1.p_2)_p| \leq 2(D_1 + \delta)$ and hence $(y_1.y_2)_p \geq (p_1.p_2)_p - 2D_1 \geq (x_1.x_2)_p - (4D_1 + 2\delta)$. Since $(y_1.y_2)_p \leq N_1$ and $(x_1.x_2)_p \geq N_1 - 1$ we have

$$|(y_1.y_2)_p - (p_1.p_2)_p| \leq (1 + 2\delta + 4D_1).$$

Next, suppose that $c_i \in [p, y_i]$, $i = 1, 2$ and $c \in [y_1, y_2]$ are the internal points of $\Delta p y_1 y_2$. We shall show that $d(p_i, c_i)$, $i = 1, 2$ are small.

Suppose $q_i \in [p, p_i]$, $i = 1, 2$ are internal points of $\triangle p p_1 p_2$. Then $d(p_i, q_i) \le d(p_1, p_2) \le D_2$. We can choose $r_i \in [p, y_i]$ such that $d(r_i, q_i) \le 2D_1$, by Lemma 1.19 applied to the subsegment of the quasi-geodesic γ_i between p, p_i and p, y_i. Hence $d(c_i, r_i) = |d(p, c_i) - d(p, r_i)| \le |d(p, c_i) - d(p, q_i)| + |d(p, q_i) - d(p, r_i)| \le (1 + 2\delta + 4D_1) + d(q_i, r_i)$. Hence $d(c_i, r_i) \le (1 + 2\delta + 6D_1)$. This gives $d(c_i, p_i) \le d(c_i, r_i) + d(r_i, q_i) + d(q_i, p_i) \le (1 + 2\delta + 8D_1 + D_2)$. Since $d(c, c_i) \le 4\delta$ we have

$$d(c, p_i) \le d(c, c_i) + d(c_i, p_i) \le (1 + 6\delta + 8D_1 + D_2).$$

Thus for any point $y_3 \in [p_1, p_2]$ we have $d(c, y_3) \le d(p_1, p_2) + d(p_1, c) \le (1 + 6\delta + 8D_1 + 2D_2) = D_3$, say.

Proof of Condition 1 : Now,

$$\begin{aligned} &\textstyle\sum_{i=1}^{2} l(\widehat{p_i y_i}) + d(p_1, p_2) \\ &\le \textstyle\sum_{i=1}^{2} \{\lambda d(p_i, y_i) + \lambda\} + D_2, \text{ since } \gamma_i \text{ are } \lambda-\text{quasi-geodesics.} \\ &\le \textstyle\sum_{i=1}^{2} \lambda\{d(y_i, c) + d(c, p_i)\} + 2\lambda + D_2 \\ &\le \lambda d(y_1, y_2) + 2\lambda + D_2 + 2\lambda D_3. \end{aligned}$$

Proof of Condition 2 :

$$\begin{aligned} &l(\widehat{p_1 y_1}) + d(p_1, y_3) \\ &\le \{\lambda d(p_1, y_1) + \lambda\} + d(p_1, y_3) \\ &\le \lambda\{d(y_1, y_3) + d(y_3, p_1)\} + \lambda + d(p_1, y_3) \\ &\le \lambda d(y_1, y_3) + \lambda + (\lambda + 1) d(p_1, p_2) \\ &\le \lambda d(y_1, y_3) + (\lambda + (\lambda + 1) D_2). \end{aligned}$$

The proof of Condition 3 is exactly like the proof of Condition 2. □

Using stability of quasi-geodesics (Lemma 1.19) the proofs of the following lemma and corollary are standard (see Lemma 1.15, Chapter *III.H*, [BH99]).

Lemma 2.5. Asymptotic rays are uniformly close: *For all $\delta \ge 0$ and $k \ge 1$ there is a constant $D = D(\delta, k)$ such that the following holds:*

Suppose X is a δ hyperbolic metric space and $\gamma_1, \gamma_2 : [0, \infty) \to X$ are two asymptotic k-quasi-geodesic rays. Then there exists $T \ge 0$ such that $\gamma_1(t) \in N_D(Im(\gamma_2))$ and $\gamma_2(t) \in N_D(Im(\gamma_1)) \le D$, for all $t \ge T$.

Corollary 2.6. *For all $\delta \ge 0$ and $K \ge 1$ there is a constant $D_{2.6} = D_{2.6}(\delta, K)$ such that the following holds:*

Suppose X is a δ-hyperbolic metric space and let γ_1, γ_2 be two K-quasi-geodesic lines in X joining the same pair of points $\xi_1, \xi_2 \in \partial X$. Then the Hausdorff distance between γ_1 and γ_2 is at most $D_{2.6}$.

Lemma 1.19 and Lemma 2.5 combined with the proof of Lemma 1.22, immediately imply the following result.

Lemma 2.7. *For all $\delta \ge 0$, $D^{'} \ge 0$ and $k \ge 1$ there are constants $D = D_{2.7}(\delta, k)$ and $L = L_{2.7}(\delta, k, D^{'})$ such that we have the following:*

Suppose X is a δ-hyperbolic metric space. Then

1. *Let $\Delta \xi_1 \xi_2 \xi_3$ be a k-quasi-geodesic* ideal triangle *in X, i.e. a union of three k-quasi-geodesic lines in X joining the pairs of points (ξ_i, ξ_j). Let us denote the quasi-geodesic lines by $[\xi_i, \xi_j]$. Then there is a point $x \in X$ such that $x \in N_D([\xi_i, \xi_j])$ for all $i \ne j$.*

(2) *If $x, x^{'} \in X$ are two points each of which is contained within a $D^{'}$- neighborhood of each of the sides of an ideal k-quasi-geodesic triangle $\Delta\xi_1\xi_2\xi_3$ then $d(x, x^{'}) \leq L$.*

If a point $x \in X$ is contained in the $D^{'}$-neighborhood of each of the sides of an ideal quasi-geodesic triangle $\Delta\xi_1\xi_2\xi_3$, then x will be called a $D^{'}$*-barycenter* of $\triangle\xi_1\xi_2\xi_3$. A $D_{2.7}$-barycenter will be simply referred to as a *barycenter.*

Now, Lemma 2.7 along with the proof of Lemma 1.22(2) gives the following.

Lemma 2.8. *Given $\delta \geq 0$, $D^{'} \geq 0$, $K_1 \geq 1$ and $K_2 \geq 0$ there exists $D = D(\delta, K_1, K_2, D^{'})$ such the following holds:*

Suppose $f : X \to Y$ is a K_1-quasi isometric embedding of δ-hyperbolic metric spaces. Let $\triangle\xi_1\xi_2\xi_3 \subset X$ and $\triangle\partial(f)(\xi_1)\partial(f)(\xi_2)\partial(f)(\xi_3) \in Y$ be K_2-quasi-geodesic ideal triangles. If $x \in X$ is a $D^{'}$-barycenter of $\Delta\xi_1\xi_2\xi_3$, then $f(x) \in Y$ is a D-barycenter of $\Delta\partial(f)(\xi_1)\partial(f)(\xi_2)\partial(f)(\xi_3)$.

The barycenter map
Suppose X is a δ-hyperbolic metric space such that ∂X has more than two points. Let us denote the set of all distinct triples of points in ∂X by $\partial^3 X$. Now, given $\xi = (\xi_1, \xi_2, \xi_3) \in \partial^3 X$ we can, by Lemma 2.4, construct a $K_{2.4}(\delta)$-quasi-geodesic ideal triangle, say Δ_1, with vertices ξ_i, $i = 1, 2, 3$. Then, by Lemma 2.7(2) there is a coarsely well defined barycenter of Δ_1. Suppose b_ξ is a barycenter of Δ_1. *Henceforth, we shall refer to it simply as a barycenter of the triple* (ξ_1, ξ_2, ξ_3). For a different set of choices of the $K_{2.4}(\delta)$-quasi-geodesic lines joining the pairs (ξ_i, ξ_j), suppose we obtain a new ideal triangle Δ_2, and suppose $b^{'}_\xi$ is a barycenter of (ξ_1, ξ_2, ξ_3) defined with respect to Δ_2. Then by the stability of quasi-geodesic lines (Lemma 2.6), $b^{'}_\xi$ is a $D_1 := (D_{2.7}(\delta) + D_{2.6}(\delta, K_{2.4}(\delta))$-barycenter of the triangle Δ_1. Hence, by Lemma 2.7(2), $d(b_\xi, b^{'}_\xi) \leq L_{2.7}(\delta, K_{2.4}(\delta), D_1)$ and we have:

Lemma 2.9. *For every $\delta \geq 0$ there is a constant $D_{2.9} = D_{2.9}(\delta)$ such that we have the following:*

Suppose X is a δ-hyperbolic metric space and $\xi = (\xi_1, \xi_2, \xi_3) \in \partial^3 X$. If b_ξ and $b^{'}_\xi$ are two barycenters of ξ, then $d(b_\xi, b^{'}_\xi) \leq D_{2.9}$.

We shall say that a map $f : (U, d_U) \to (V, d_V)$ satisfying properties $\mathcal{P}_1, \cdots, \mathcal{P}_k$ is *coarsely unique* if there exists $C > 0$ such that for any other map $g : (U, d_U) \to (V, d_V)$ satisfying properties $\mathcal{P}_1, \cdots, \mathcal{P}_k$, and any $u \in U$, $d_V(f(u), g(u)) \leq C$.

Thus, from Lemma 2.9 we have a coarsely unique map $\phi : \partial^3 X \to X$, $\xi \mapsto b_\xi$ mapping a triple of points to a barycenter. Any such map will be referred to as *the barycenter map.*

Recall that a family of maps $f_\alpha : X_\alpha \to Y_\alpha$, from sets X_α to metric spaces (Y_α, d_{Y_α}), α running over some indexing set $\mathcal{F}$, is said to be *uniformly coarsely surjective* if there is a constant $D \geq 0$, such that for all $\alpha \in \mathcal{F}$, Y_α is contained in the D-neighborhood of $f_\alpha(X_\alpha)$.

Now we are ready to state the main proposition of this subsection. For any graph X we will denote its vertex set by $\mathcal{V}(X)$.

Proposition 2.10. Existence of qi sections for metric graph bundles: *For all $\delta^{'}, N \geq 0$ and proper $f : \mathbb{R}^+ \to \mathbb{R}^+$ there exists $K_0 = K_0(f, \delta^{'}, N)$ such that the following holds.*
Suppose $p : X \to B$ is an (f, K)-metric graph bundle with the following properties:

(1) *Each of the fibers* F_b , $b \in \mathcal{V}(B)$ *is a* δ'*-hyperbolic metric space with respect to the path metric* d_b *induced from* X*.*
(2) *The barycenter maps* $\phi_b : \partial^3 F_b \to F_b$ *are uniformly coarsely surjective, i.e. for all* $b \in B$*, the image of* ϕ_b *is contained in* $\mathcal{V}(F_b)$ *and* F_b *is contained in the* N*-neighborhood of the image of* ϕ_b*.*

Then there is a K_0*-qi section through each point of* $\mathcal{V}(X)$*.*

Note that the constant K in Proposition 2.10 above is given by $K = f(4)$ by Proposition 1.4 and hence we may write $K_0 = K_0(f, K, \delta', N)$ by making the implicit dependence on K explicit.

Before we start the proof a remark is in order. By a qi section of a metric graph bundle we will always mean, from now on, a section defined only on the vertex set $\mathcal{V}(B)$ with image contained in $\mathcal{V}(X)$. Such a qi section can be easily extended over all of B by mapping edges of B linearly over edge-paths in X of uniformly bounded length.

Proof. Let us fix a set $\{\phi_b\}_{b\in\mathcal{V}(B)}$ of barycenter maps and let $v \in \mathcal{V}(B)$, $x \in \mathcal{V}(F_v)$. First, suppose that x is contained in the image of the barycenter map ϕ_v. We will construct a qi section through x. Choose a point $\xi_v = (\xi_1, \xi_2, \xi_3) \in \partial^3 F_v$ such that $\phi_v(\xi_v) = x$. Let $w, z \in \mathcal{V}(B)$, $w \neq z$. Choose a geodesic γ joining w, z and let $w = w_0, w_1, \ldots, w_{n-1}, w_n = z$ be the consecutive vertices on γ. By condition (2)(ii) of the definition of metric graph bundles (Proposition 1.4), for all i, $0 \le i \le n-1$, there is a K-quasi-isometry $f_{w_i w_{i+1}} : F_{w_i} \to F_{w_{i+1}}$ which sends any vertex $y_i \in \mathcal{V}(F_{w_i})$ to a vertex $y_{i+1} \in \mathcal{V}(F_{w_{i+1}})$ where y_i and y_{i+1} are connected by an edge. By composition of n such maps we get a map $f_{wz} : F_w \to F_z$, which sends each point $y \in F_w$ to a point $y' \in \mathcal{V}(F_z)$ such that $d(y, y') \le n = d_B(w, z)$. Let $f_{ww} : F_w \to F_w$ denote the identity map, for all $w \in \mathcal{V}(B)$. Now we make the following observations:

1. Since the inclusion maps $F_w \hookrightarrow X$ are uniformly metrically proper, by the definition of metric graph bundles, the map f_{wz} is coarsely uniquely determined. In fact, if $d(w, z) = n$, $n \in \mathbb{N}$, we have for any other map f'_{wz} defined in the same way, $d(f_{wz}(y), f'_{wz}(y)) \le 2n$, so that $d_z(f_{wz}(y), f'_{wz}(y)) \le f(2n)$, for all $y \in F_w$.

2. Since each f_{wz} is obtained as a composition of K-quasi-isometries it is a quasi-isometry. Now, since the spaces $F_w, w \in \mathcal{V}(B)$, are δ'-hyperbolic and since the map f_{wz} is coarsely uniquely determined, we have a well defined map $\partial(f_{wz}) : \partial F_w \to \partial F_z$, by Lemma 2.2, and hence an induced map $\partial^3 f_{wz} : \partial^3 F_w \to \partial^3 F_z$, $\forall w, z \in \mathcal{V}(B)$.

Denote $\xi_v = \xi$ and $\phi_v(\xi) = x$. Consider the map $s = s_{\xi,x} : \mathcal{V}(B) \to X$ given by $s(v) = x = \phi_v(\xi)$, and $s(w) = \phi_w((\partial^3 f_{vw}(\xi)))$, for all $w \in \mathcal{V}(B), w \neq v$. We show below that s (or $s(\mathcal{V}(B))$) is the required qi section through x.

3. Writing $\xi_w = \partial^3 f_{vw}(\xi)$ for all $w \neq v$, note that for any $w, z \in \mathcal{V}(B)$, $\partial^3 f_{wz}(\xi_w) = \xi_z$.
This follows from the fact that by the definitions of the maps f_{vz}, f_{wz}, f_{vw} we have, for all $y \in F_v$, $d(f_{vz}(y), f_{wz} \circ f_{vw}(y)) \le d_B(v, z) + d_B(w, z) + d_B(v, w) + 3$, and thus $d_z(f_{vz}(x), f_{wz} \circ f_{vw}(x)) \le f(d_B(v, z) + d_B(w, z) + d_B(v, w) + 3)$. The claim follows from Lemma 2.2(3).

4. Lastly, if w, z are adjacent vertices of B then $d(s(w), s(z)) \le C$ for some constant C. This can be shown as follows. If $w, z \in \mathcal{V}(B)$ are adjacent then we know that the map f_{wz} is a K-quasi-isometry. Let $\xi_w = (\beta_1, \beta_2, \beta_3)$ and $\partial^3 f_{wz}(\xi_w) = \xi_z =$

(η_1, η_2, η_3). Choose $K_{2.4}(\delta^{'})$-quasigeodesic ideal triangles Δ_w and Δ_z respectively in F_w and F_z, with vertices ξ_i's and η_i's; by definition of the map s, $s(w)$ and $s(z)$ are $D_{2.7}(\delta^{'})$-barycenters of these triangles. Now, the map f_{wz} takes the ideal triangle Δ_w to an ideal K_1-quasigeodesic triangle with vertices η_i's, where $K_1 = K_{2.4}(\delta^{'}).K + K$, and $f_{wz}(s(w))$ is a $D_1 := \{D_{2.7}(\delta^{'}).K + K\}$-barycenter of the new triangle. Thus, by Lemma 2.6, $f_{wz}(s(w))$ is a D_2-barycenter of the triangle Δ_z, where $D_2 = D_{2.6}(\delta^{'}, K_1) + D_1$. Hence, by Lemma 2.7, we have $d(s(z), f_{wz}(s(w))) \leq L_{2.7}(\delta^{'}, K_{2.4}(\delta^{'}), D_2)$. Since $d(s(w), f_{wz}(s(w))) = 1$ we have $d(s(w), s(z)) \leq C := 1 + L_{2.7}(\delta^{'}, K_{2.4}(\delta^{'}), D_2)$.

Now it is clear that s is a C-qi section over $\mathcal{V}(B)$: For $w, z \in \mathcal{V}(B)$, we have $d(w, z) \leq d(s(w), s(z))$ by the definition of a metric graph bundle. Also from Step (4) above, we have $d(s(w), s(z)) \leq C.d(w, z) + C$.

If $x \in \mathcal{V}(F_v)$ is not in the image of ϕ_v, we can choose $x_1 \in \mathcal{V}(F_v)$ such that $d(x, x_1) \leq N$ and $x_1 \in Im(\phi_v)$. Now construct as above a C-qi section $s = s_{\xi, x_1}$, and define a new section $s^{'}$ by setting $s^{'}(b) = s(b)$ for all $b \in \mathcal{V}(B), b \neq v$ and $s^{'}(v) = x$. This is an $(N + C)$-qi section passing through x. Thus we can take $K_0 = N + C$ to finish the proof of the proposition. □

Applying this proposition to Example 1.5, we have a different proof of the following result of Mosher [Mos96].

Theorem 2.11. (Mosher [Mos96]**)** *Let us consider the short exact sequence of finitely generated groups*

$$1 \to A \to G \to Q \to 1.$$

such that A is non-elementary word hyperbolic. Then there exists a q(uasi)-i(sometric) section $\sigma : Q \to G$*. Hence, if G is hyperbolic, then so is Q.*

Remark 2.12. *We note first that if the fibers of a metric bundle are (uniformly) hyperbolic, then so are the vertex spaces of an approximating metric graph bundle. This is because the fibers of an approximating metric graph bundle are uniformly quasi-isometric to the fibers of the metric bundle.*

Next (for the same reason) observe that if the barycenter maps of the metric bundle are uniformly coarsely surjective, then so are the barycenter maps of an approximating metric graph bundle.

It is not hard to see using Remark 2.12 that if the horizontal spaces of a metric bundle X (as opposed to a metric graph bundle) satisfy the two conditions of Proposition 2.10 then through each point of X there is a qi section. One needs only a slight modification of the proof of Proposition 2.10. However, we shall give a simple proof below using Proposition 2.10.

Proposition 2.13. Existence of qi section for metric bundles: *Let* $p : X^{'} \to B^{'}$ *be an* (f, c, K)*-metric bundle and let* $\pi : X \to B$ *be an approximating metric graph bundle as in Lemma 1.14. Let* $\psi_B, \psi_X, \psi_{X'}, \psi_{B'}$ *denote the maps described in Lemma 1.14. Suppose these are all* K_1*-quasi-isometries. Let* $V \subset B^{'}$ *be the vertex set of B. Suppose we have a k-qi section* $s : V \to X$*. Then we have a* $k^{'} = K_{2.13}(f, c, K, K_1, k)$*-qi section* $s^{'} : B^{'} \to X^{'}$ *such that* $\psi_X \circ s = s^{'} \circ \psi_B$*.*

Thus any metric bundle satisfying the properties
1) horizontal spaces are uniformly hyperbolic, and

2) the barycenter maps of these spaces are uniformly coarsely surjective admits a uniform qi section through each point.

Proof. The proof of the first part of the proposition is clear once we describe what the map $s^{'}$ is. For $u \in V$ define $s^{'}(u) = \psi_{B'} \circ s(u)$. Suppose $u \in B \setminus V$. Let $v \in V$, so that $d(u, v) \leq 1$. Choose $x \in F_v$ such that x can be joined to $s^{'}(u)$ by a curve in $X^{'}$ of length at most c and define $s^{'}(v) = x$.

The last part of the proposition follows from Proposition 2.10, Remark 2.12 and the first part of the proposition. ☐

2.2. **Nearby qi sections.** The next lemma leads to one of the main tools (Lemma 2.17) for proving our theorem. This lemma originally appears in [Ham05] in the context of metric fibrations. The proof that we give here is almost the same as that of [Ham05], nevertheless we include it for the sake of completeness.

Definition 2.14. *Suppose X is an (f, c, K)-metric bundle (resp. (f, K)-metric graph bundle) over B. For two qi sections X_1, X_2 and $A > 0$ we define $U_A(X_1, X_2)$ to be the set $\{b \in B : d_b(X_1 \cap F_b, X_2 \cap F_b) \leq A\}$ (resp. $\{b \in \mathcal{V}(B) : d_b(X_1 \cap F_b, X_2 \cap F_b) \leq A\}$).*

Suppose X is an (f, c, K)-metric bundle over B satisfying (M_k, λ_k, n_k)-flaring for all $k \geq 1$ (cf. Definition 1.9), and let μ_k be the bounded flaring function (cf. Corollary 1.12). Then we have the following:

Lemma 2.15. *For all $c_1 \geq 1$ and $R \geq 0$ there are constants $D_{2.15} = D_{2.15}(c_1, R)$ and $K_{2.15} = K_{2.15}(c_1)$ such that the following holds.*

Suppose X_1, X_2 are two c_1-qi sections of B in X and let $A \geq M_{c_1}$.

1. *Let $\gamma : [t_0, t_1] \to B$ be a geodesic such that*
   a) $d_{\gamma(t_0)}(X_1 \cap F_{\gamma(t_0)}, X_2 \cap F_{\gamma(t_0)}) = AR$.
   b) $\gamma(t_1) \in U_A := U_A(X_1, X_2)$ but for all $t \in [t_0, t_1)$, $\gamma(t) \notin U_A$.
   Then the length of γ is at most $D_{2.15}(c_1, R)$.
2. *U_A is $K_{2.15}$-quasi-convex in B.*

For convenience of exposition we will write λ for λ_{c_1}, n for n_{c_1} and μ for μ_{c_1} in the proof of the lemma below. Also $l(\alpha)$ will denote the length of a curve α.

Proof. (1) Let $\phi : [t_0, t_1] \to \mathbb{R}$ be the function $t \mapsto d_{\gamma(t)}(X_1 \cap F_{\gamma(t)}, X_2 \cap F_{\gamma(t)})$ and $t_1 - t_0 = n.L + \epsilon$ where $L \in \mathbb{Z}^+$ and $0 \leq \epsilon < n$. Suppose we have $L \geq 3$. Now consider the sequence of numbers $\phi(t_0 + ni)$, $i = 1, \cdots, L$. Since $\phi(t_0 + n.i) \geq M_{c_1}$, for all $i \in [1, L - 1]$ we have

$$\lambda\phi(t_0 + ni) \leq \max\{\phi(t_0 + n(i - 1)), \phi(t_0 + n(i + 1))\}$$

by the flaring condition.

Thus, if $\phi(t_0 + n) > \phi(t_0)$ then $\phi(t_0 + n(i + 1)) \geq \lambda\phi(t_0 + ni)$ for all $i \in [1, L - 1]$. Then, it follows that $\phi(t_0 + nL)) \geq \lambda^{L-1}\phi(t_0)$. Now using bounded flaring (Corollary 1.12) we have, $\phi(t_0 + nL) \leq \mu(n) \max\{\phi(t_1), 1\}$. Putting all these together and using the fact that $\phi(t_1) \leq A$ and $\phi(t_0) > A$, we have $L - 1 < log(\mu(n))/log\lambda$.

Thus, $L \geq 3$ and $\phi(t_0 + n) > \phi(t_0)$ implies

$$l(\gamma) < n(L + 1) \leq 2n + n.log\mu(n)/log\lambda.$$

Now, suppose $\phi(t_0) \geq \phi(t_0 + n)$ and let $k \leq L$ be the largest integer such that $\phi(t_0) \geq \phi(t_0 + n) \geq \cdots \geq \phi(t_0 + k.n)$. If $k \geq 2$, applying the flaring condition we get

$\phi(t_0+(i-1)n) \geq \lambda\phi(t_0+in)$ for all $i \in [1, k-1]$. Then $\phi(t_0) \geq \lambda^{k-1}\phi(t_0+(k-1)n) > \lambda^{k-1}A$. Thus, in this case, we always have $k < 1 + \{log\phi(t_0) - logA\}/log\lambda = 1 + logR/log\lambda$. Also, by the first part of the proof, we know that $l(\gamma|_{[t_0+k.n,t_1]}) < \max\{3n, 2n + n\ log\mu(n)/log\lambda\}$. Hence, in any case, we have

$$l(\gamma) \quad \leq \quad n + n\{logR/log\lambda + max\{3n, 2n + n\ log(\mu(n))/log\lambda\}.$$

Taking $D_{2.15} = D_{2.15}(c_1, R)$ be the right hand side of the above inequality, part (1) of the lemma is proved.

(2) Suppose $\gamma : [t_0, t_1] \rightarrow B$ is a geodesic joining two points of U_A, such that for all $t \in (t_0, t_1)$, $\gamma(t) \notin U_A$. Without loss of generality, we may assume that $t_1 - t_0 > n$. Let $t_2 = t_0 + n$. Then by bounded flaring, we have $\phi(t_2) \leq \mu(n)\phi(t_0) \leq \mu(n).A$. Again by the first part of the lemma $l(\gamma|_{[t_2,t_1]}) \leq D_{2.15}(c_1, \phi(t_2)/A)$. Since, the function $D_{2.15}$ is increasing in the second co-ordinate, given that the first co-ordinate is fixed, we have $l(\gamma) \leq n + D_{2.15}(c_1, \mu(n))$. Hence, taking $K_{2.15}(c_1) = n + D_{2.15}(c_1, \mu(n))$ completes the proof of the lemma. □

Remark 2.16. If a metric graph bundle satisfies a flaring condition then the same results hold. The proof is the same. We note in particular that in Lemma 2.15 (1), all that we need in order to make an analogous statement for a metric graph bundle is that $\phi(t) \geq M_{c_1}$, for all t.

We indicate below what remains true for an approximating metric graph bundle of a metric bundle. Thus, Lemma 2.17 below addresses the issue that if a metric bundle satisfies the flaring condition, then the approximating metric graph bundle may not necessarily satisfy a flaring condition *per se.* However, as the lemma below shows, in spite of this, conclusions similar to those of Lemma 2.15 above remain true. These conclusions are true for a metric graph bundle satisfying a flaring condition as noted earlier. In the next two sections where we give a proof of the main theorem 4.10 we replace a metric bundle with flaring by its approximating metric graph bundle. Then we work exclusively with a metric graph bundle. Thus the conclusion of the lemma below can be used both for a metric graph bundle satisfying a flaring condition as also the approximating metric graph bundle of a metric bundle that satisfies a flaring condition.

Let $p : X^{'} \rightarrow B^{'}$ be an (f, c, K)-metric bundle satisfying (M_k, λ_k, n_k)-flaring for all $k \geq 1$ and let $\pi : X \rightarrow B$ be an approximating metric graph bundle as in Lemma 1.14. As in the remark after Lemma 1.14 we suppose that all the maps $\psi_X, \psi_B, \psi_{X'}, \psi_{B'}$, as also the restrictions of ψ_X and $\psi_{X'}$ to horizontal spaces, are K_1- quasi-isometries. Let B be λ-hyperbolic. Suppose further that for every k-qi section of the approximating metric graph bundle, we obtain a $k^{'}$-qi section of the original bundle. For convenience of exposition we suppress the dependence of the constants (defined in the following lemma) on the parameters f, c, K, λ etc.

Lemma 2.17. *With notation as above, let X_1, X_2 be two k-qi sections of the approximating metric graph bundle. Suppose $d_b(F_b \cap X_1, F_b \cap X_2) \geq A_0 \geq 0$ for all $b \in \mathcal{V}(B)$ and let $A_1 = K_1.max\{A_0 + K_1 + 1, M_{k'} + K_1\}$. Then the following hold.*
(1) For $A \geq A_1$, $U_A(X_1, X_2)$ is $K_{2.17} = K_{2.17}(k, A)$-quasi-convex in B.
(2) Suppose $d_u(F_u \cap X_1, F_u \cap X_2) = C \geq A$ for some $u \in \mathcal{V}(B)$. Then $d(u, U_A(X_1, X_2)) \leq D_{2.17}(k, C)$.

(3) *Suppose* $A_0 \geq K_1(M_{k'} + K_1)$. *Then the diameter of the set* $U_A(X_1, X_2)$ *is at most* $D'_{2.17}(k, A)$.

Proof. For the proof of this lemma we introduce the notation d'_u for the path metric of X'_u induced from X'. Also let $A_2 = A_1/K_1 - 1$.
(1) By Proposition 2.13 we have two k'-qi sections X'_1, X'_2 of B' in X' (corresponding to X_1, X_2 respectively) where $k' = K_{2.13}(f, c, K, K_1, k)$. By choice of the constant A_1, we know that $U := U_{A_2}(X'_1, X'_2)$ is a nonempty $K' := K_{2.15}(k')-$quasiconvex subset of B'. Hence $\psi_{B'}(U) \subset B$ is $D := \{K_1.K' + K_1 + D_{1.19}(\lambda, K_1)\}$-quasiconvex. Also note that $\psi_{B'}(U) \subset U_A(X_1, X_2)$.

Now suppose $u \in (U_A(X_1, X_2) \setminus \psi_{B'}(U))$ is a point of $\mathcal{V}(B)$. It follows that $d'_u(X'_1 \cap F_u, X'_2 \cap F_u) \geq M_{k'}$; else $u \in U$ would imply that $u = \psi_{B'}(u) \in \psi_{B'}(U)$.

Hence from Lemma 2.15(1), $d_{B'}(u, U) \leq D_1 = D_{2.15}(k', (A.K_1 + K_1)/A_2)$. Therefore $d_B(u, \psi_{B'}(U)) \leq D_2 = K_1 D_1 + K_1$. Thus $\psi_{B'}(U) \subset U_A(X_1, X_2) \subset N_{D_2}(\psi_{B'}(U))$. Since $\psi_{B'}(U)$ is a D-quasi-convex set and since B is λ-hyperbolic, it follows that $U_A(X_1, X_2)$ is $K_{2.17}(= (2\lambda + D + D_2))-$quasi-convex.

(2) If $u \in U$ then we set $D_{2.17}(k, C) = 0$. Otherwise $d'_u(X'_1 \cap F_u, X'_2 \cap F_u) \leq C.K_1 + K_1$ since the restriction of the map ψ_X to the horizontal space F_u is a K_1-quasi-isometry. Hence by Lemma 2.15(1) we have $d_{B'}(u, U) \leq D_3 = D_{2.15}(k', (C.K_1 + K_1)/A_2)$. Using the fact that $\psi_{B'}$ is a K_1-quasi-isometry, we have $d_B(u, \psi_{B'}(U)) \leq K_1.D_3 + K_1$. Hence $d(u, U_A(X_1, X_2)) \leq K_1 D_3 + K_1$. Set $D_{2.17}(k, C) = K_1 + K_1.D_{2.15}(k', A_3)$ where $A_3 = max\{1, (C.K_1 + K_1)/A_2\}$.

(3) By the given condition, for all $z \in U$, $d'_z(X'_1 \cap F_z, X'_2 \cap F_z) \geq M_{k'}$ and so we can apply the flaring condition. Now let $b_1, b_2 \in U$ and let $b \in [b_1, b_2]$; then $d'_b(X'_1 \cap F_b, X'_2 \cap F_b) \leq \mu_{k'}(K_{2.15}(k')) = D_4$, say, by Corollary 1.12.

Finally, as noted in Remark 2.16, what we really used in the proof of Lemma 2.15(1) is the fact that the value of the function ϕ is always greater than or equal to $M_{k'}$. Thus in the same way we have $d_{B'}(b_1, b_2) \leq D_{2.15}(k', D_4/A_2)$. Taking $D'_{2.17}(k, A) = K_1 + K_1.D_{2.15}(k', D_4/A_2)$ completes the proof of the lemma. ☐

3. Construction of Hyperbolic Ladders

In this section we prove the main technical result leading to the combination theorem 4.10. A brief sketch follows. For a metric bundle, we first replace it with its approximating metric graph bundle. The rest of the proof deals with metric graph bundles. Then, we define *ladders* in a metric graph bundle, which are analogs of 'hallways' in [BF92] and the 'ladders' constructed in [Mit98b] and [Mit98a]. These are coarse I-bundles over the base space of the metric graph bundle. In section 3.2 we prove that ladders are hyperbolic metric spaces under the assumption that the original metric (graph) bundle satisfies a flaring condition. To achieve this we first prove this result in section 3.1 when the ladder is of small girth (i.e. the I-bundle has fibers of small enough length at some point of the base space, see Definition 3.2). Then, to prove hyperbolicity in the general case, a ladder is decomposed into small-girth ladders using qi sections. In this way, we obtain a finite sequence of hyperbolic metric spaces and check that the conditions of Corollary 1.42 are satisfied. This completes the sketch of the proof.

Notation and conventions: For the next two subsections, we fix the following notation and conventions. For us $p : X \to B$ will be either an $f-$ metric graph bundle satisfying a flaring condition, or an approximating $(f-)$ metric graph bundle obtained from a metric bundle $\pi : X^{'} \to B^{'}$ satisfying a flaring condition as in Lemma 1.14. In the latter case, we shall assume that all the maps ψ_B, ψ_X etc and their restriction to the fiber spaces are K_1-quasi-isometries.

The symbols g, μ_k will have the same connotation as in Lemma 1.10 and Corollary 1.12 respectively. We shall assume that B is δ-hyperbolic and each of the horizontal spaces F_b is $\delta^{'}$-hyperbolic for all vertices b of B. We assume that the barycenter maps $\partial^3 F_b \to F_b$ are (uniformly) coarsely surjective. Thus by Proposition 2.10 we know that the metric graph bundle admits uniform (K_0, say)qi sections through each point of X. Lastly, often the dependence on these functions and constants will not be explicitly stated if it is clear from context. The vertex set of a graph Y will be denoted by $\mathcal{V}(Y)$ as usual. By points in a graph we shall always mean vertices, unless otherwise specified.

We use the term *ladder* below due to a similar ladder construction in [Mit98b]. The term *girth* is taken from [BF92].

Definition 3.1. *Suppose X_1 and X_2 are two c_1-qi sections of the metric graph bundle X, over $\mathcal{V}(B)$. For each $b \in \mathcal{V}(B)$, join the points $X_1 \cap F_b$, $X_2 \cap F_b$ by a geodesic in F_b. We denote the union of these geodesics by $C(X_1, X_2)$, and call it a* **ladder** *formed by the sections X_1 and X_2.*

Since the Hausdorff distance between any pair of ladders determined by two given sections X_1, X_2 is uniformly bounded, $C(X_1, X_2)$ will refer to any one of them, and abusing notation we refer to it as *the* ladder determined by X_1, X_2.

For four qi sections X_i, $i = 1, 2, 3, 4$ we write $C(X_3, X_4) \subset C(X_1, X_2)$ to mean that over each point of the base space the horizontal geodesic of $C(X_3, X_4)$ is contained in the horizontal geodesic of $C(X_1, X_2)$.

Definition 3.2. *Suppose X_1 and X_2 are two c_1-qi sections. We define $d_h(X_1, X_2)$ to be the quantity* $\inf_b\{d_b(F_b \cap X_1, F_b \cap X_2) : b \in \mathcal{V}(B)\}$ *and call it the* **girth** *of the ladder $C(X_1, X_2)$.*

Lemma 3.3. *For all $c_1 \geq 1$, there exists $C_{3.3}(= C_{3.3}(c_1))$ such that the following holds.*
Suppose X_1 and X_2 are two c_1-qi sections. Then through each point $x \in C(X_1, X_2)$ there exists a $C_{3.3}$-qi section contained in $C(X_1, X_2)$.

Proof. We already know that there is a K_0-qi section, say Y_1, through x in X. Now define a new section Y_2 as follows: let $Y_2 \cap F_b$ be a nearest point projection (in the intrinsic metric on the horizontal space F_b) of $Y_1 \cap F_b$ onto the horizontal geodesic $C(X_1, X_2) \cap F_b$. This defines a set theoretic section. We need to check that this is indeed a qi section. For this it is enough to check that $\forall\, b_1, b_2 \in \mathcal{V}(B)$, with $d(b_1, b_2) = 1$, the distance between $F_{b_1} \cap Y_2$ and $F_{b_2} \cap Y_2$ is uniformly bounded. This follows immediately from Lemma 1.10, and Lemma 1.27 by choosing $C_{3.3} := c' + D_{1.27}(\delta', g(c'))$, where $c' = 2\ max\{K_0, c_1\}$. $\square$

Simplification of Notation: We fix the following conventions and notation to be followed in the rest of this section. Fix $c_1 \geq K_0$. Let $c_{i+1} = C^i_{3.3}(c_1)$, $i = 1, 2, 3$ where $C^i_{3.3}$ is the i-th iterate of the function $C_{3.3}$. Note that if Y is a k-qi section, and $k \leq c_4$, then it is also a c_4-qi section. We know that our metric graph bundle satisfies the bounded flaring condition. We shall denote the function μ_{c_4} (see Corollary 1.12) simply by μ.

With reference to Proposition 2.13 we assume that any c_i qi section of the approximating metric graph bundle gives rise to a $c_i^{'}$-qi section of the original metric bundle.

Remark 3.4. (1) From the definition of a ladder, we see that a ladder in a metric graph bundle is *not* connected; but for any $D \geq 2c_1$, the D-neighborhood of a ladder formed by two c_1-qi sections is indeed connected. In general, we shall work with a large enough neighborhood of a ladder, and since we are dealing with graphs, these neighborhoods are then geodesic metric spaces.

(2) Also note that a c_2-qi section lying inside a ladder formed by two c_1-qi sections is in fact the image of a $2c_2$-Lipschitz map from $\mathcal{V}(B)$ into the D-neighborhood ($D \geq 2c_2$) of the ladder equipped with the path metric induced from X. Since the projection map of the metric graph bundle X to its base space B is 1-Lipschitz by definition, it follows that a c_2-qi section lying inside a ladder formed by two c_1-qi sections is a $2c_2$-qi section inside a large enough neighborhood of the ladder.

(3) By paths in X or B we always mean a sequence of vertices. Suppose $b_0, b_1, \ldots, b_n$, is the sequence of vertices on a geodesic edge path γ of B. By a lift of γ in a k-qi section X_1, we mean the discrete path $\tilde{\gamma}$ obtained from the lifts $x_i \in F_{b_i} \cap X_1$, of the vertices b_i, $1 \leq i \leq n$ in X_1. It follows that $l(\tilde{\gamma}) \leq 2k.l(\gamma)$.

(4) We remarked earlier that Lemma 2.17 holds for a metric graph bundle with a flaring condition as well. Hence in what follows we use it without any further explanation.

We shall later use another construction of [Mit98a]. Let X_1, X_2 be two c_1-qi-sections of a metric graph bundle $p : X \to B$, where each fiber (but not necessarily the base B) is uniformly δ-hyperbolic. Let $C(X_1, X_2)$ be the associated ladder. By construction, the intersection $C(X_1, X_2) \cap F_b$ of the ladder with a fiber F_b is a geodesic $\lambda_b \subset F_b$ in the path metric space (F_b, d_b). Define a projection $\pi_b : F_b \to \lambda_b$ to be the nearest point projection of F_b onto the geodesic λ_b in the metric d_b and let $\Pi_{X_1,X_2} : X \to C(X_1, X_2)$ be given by

$$\Pi_{X_1,X_2}(x) = \pi_b(x),\ \forall x \in F_b.$$

The main technical Theorem of [Mit98a] states

Theorem 3.5. [Mit98a] *For X, B, p as above, and $c_1 \geq 1$, there exists $C \geq 1$ such that for two c_1-qi sections X_1, X_2, and $\forall x, y \in X$,*

$$d(\Pi_{X_1,X_2}(x), \Pi_{X_1,X_2}(y)) \leq Cd(x, y) + C.$$

Equivalently, Π_{X_1,X_2} is a coarse Lipschitz retract of X onto $C(X_1, X_2)$.

The most important ingredients, for the proof of the results that follow in this section, are Lemma 1.10, Corollary 1.12, Lemma 2.17 and Lemma 3.3.

3.1. **Hyperbolicity of ladders: Special case.**

This subsection is devoted to proving the hyperbolicity of small girth ladders.

Let X_1, X_2 be two c_1-qi sections in X and let $d_h(X_1, X_2) = A_0$, say. Let $A := K_1\ max\{A_0 + K_1 + 1, M_{c_4^{'}} + K_1\}$ (K_1 refers to notation fixed at the beginning of Section 3 and $M_{c_4^{'}}$ refers to the constant in the flaring condition). We further assume, with reference to Lemma 2.17 that for any two k-qi sections X_3, X_4, $k \leq c_4$, lying inside the ladder $C(X_1, X_2)$, the set $U_A(X_3, X_4) \subset B$ is K-quasi-convex. We shall write simply $U(X_3, X_4)$ instead of $U_A(X_3, X_4)$ in what follows. Dependence of constants in the various lemmas and propositions below on the constants associated with the bundle will be implicit rather than explicit.

The rest of this subsection is devoted to proving the following:

Proposition 3.6. *For all $L \geq 2c_4$, and c_1, A_0 as above, there exist $\delta_{3.6}(= \delta_{3.6}(c_1, A_0, L)) \geq 0$, $K_{3.6}(= K_{3.6}(c_1, A_0, L)) \geq 0$, $D_{3.6}(= D_{3.6}(c_1, A_0, L)) \geq 0$ such that we have the following:*
(1)The L-neighborhood of the ladder $C(X_1, X_2)$ in X is $\delta_{3.6}$-hyperbolic with the path metric induced from X, and X_1,X_2 are $K_{3.6}$-quasiconvex in $C(X_1, X_2)$.
(2) If $d_h(X_1, X_2) \geq K_1.M_{c_1^{'}} + K_1^2$, then X_1, X_2 are $D_{3.6}$-cobounded in the L-neighborhood of $C(X_1, X_2)$.

Idea of the proof: The proof of this proposition is rather long. Therefore, we shall break it up into several lemmas. The idea is as follows. We define a set of curves $c(x, y)$, one for each pair of points $x, y \in C(X_1, X_2)$ and check that they satisfy the three properties of Corollary 1.29. Given x, y in the ladder, first we construct two qi sections through them. Then, $c(x, y)$ consists of three parts: two of them are in the two sections containing x, y and the other one is a horizontal geodesic of uniformly bounded length. Then any problem of length computation is transferred to the sections. For instance computing the Hausdorff distance between two curves or proving slimness of triangles becomes easy when we apply this strategy to the parts of the curves that already lie in a quasi-isometric section of the hyperbolic base space B. Lemma 2.17 and the bounded flaring condition are the main tools of the proof.

We denote by $d^{'}$ the path metric on a ladder induced from X. Also $Hd^{'}$ will denote the Hausdorff distance between sets in a ladder and Hd_B will denote Hausdorff distance between sets in B.

Definition of curve family: Let $x, y \in C(X_1, X_2)$ be two vertices. By Lemma 3.3 we can choose two c_2-qi sections X_3 and X_4 through x and y respectively (see Remark 3.4 (1)), in $C(X_1, X_2)$. Recall that $U(X_3, X_4) \subseteq B$ is a K-quasi-convex subset. Join $p(x)$ to $U(X_3, X_4)$ by a shortest geodesic $\gamma_{x,y}$ in B ending at $b_{x,y} \in U(X_3, X_4)$. Let $\tilde{\gamma}_{x,y}$ be the lift of $\gamma_{x,y}$ in X_3, ending at $s_{x,y}$. Let $t_{x,y}$ be the lift of $b_{x,y}$ in X_4. We note that $d_{b_{x,y}}(t_{x,y}, s_{x,y}) \leq A$. Now let $\beta_{x,y}$ be a geodesic in B joining $p(y)$ and $b_{x,y}$, and let $\tilde{\beta}_{x,y}$ be the lift of $\beta_{x,y}$ in X_4. We define $c(x, y)$ to be the union of the three curve segments: $\tilde{\gamma}_{x,y}$, $\tilde{\beta}_{x,y}$ and the segment of $F_{b_{x,y}} \cap C(X_1, X_2)$ between $t_{x,y}$ and $s_{x,y}$. We see that there is an asymmetry in the definition of $c(x, y)$ and a number of choices are involved. However, for each unordered pair $\{x, y\}$ make the choices once and for all and choose either $c(x, y)$ or $c(y, x)$ as the curve joining the points x, y. Let us denote this set of curves by $\mathcal{S}$. (See figure below.)



Path families: Special case

Lemma 3.7. *Given $D_1 \geq 0$ and c_1, A as above, there exist constants $D_{3.7} = D_{3.7}(c_1, A, D_1)$ and $D'_{3.7} = D'_{3.7}(c_1, A, D_1)$ such that the following holds:*

Let $x, y \in C(X_1, X_2)$ with $d(x, y) \leq D_1$. Then, $d'(x, y)$- the distance between x, y in the metric of the ladder $C(X_1, X_2)$, is bounded by $D'_{3.7}$. Moreover, the length of the curve $c(x, y) \in \mathcal{S}$ is at most $D_{3.7}$.

Proof. Let $\tilde{y}$ be the lift of $p(x)$ in X_4. Since p is a 1-Lipschitz map, $d_B(p(y), p(\tilde{y})) \leq D_1$. Hence, $d(y, \tilde{y}) \leq c_2.D_1 + c_2$, since X_4 is a c_2-qi section. Hence $d(\tilde{y}, x) \leq c_2.D_1 + D_1 + c_2$. Then, since fibers of the map p are uniformly properly embedded with properness function f, we have $d_{p(x)}(\tilde{y}, x) \leq f(c_2.D_1 + D_1 + c_2)$.

By Remark 3.4(3), $d'(y, \tilde{y}) \leq 2c_2.D_1$. Thus $d'(x, y) \leq 2c_2.D_1 + f(c_2.D_1 + D_1 + c_2)$ and the first part of the lemma is proved, with $D'_{3.7} := 2c_2.D_1 + f(c_2.D_1 + D_1 + c_2)$.

Next by Lemma 2.17(2), we have

$$d_B(p(x), b_{x,y}) \leq D'_1 := D_{2.17}(c_2, f(c_2.D_1 + D_1 + c_2)).$$

Thus $d_B(p(y), b_{x,y}) \leq D_1 + D'_1$. From this and Remark 3.4(3), the second part of the lemma follows immediately, with $D_{3.7} := A + 2c_2.D_1 + 4c_2.D'_1$. □

Remark 3.8. Note that in Lemma 3.7, we have *not* assumed that $C(X_1, X_2)$ is of small girth.

We next show that the curve family is **coarsely well-defined**, i.e. ambiguities in the definition of the curves can be ignored. More precisely, the different choices of curves joining the same pair of points are at a uniformly bounded Hausdorff distance from each other.

Suppose X_3, X'_3 are two k-qi sections in $C(X_1, X_2)$ containing x; and X_4 is a k-qi section containing y, where $k \leq c_4$. Consider the two curves $c(x, y)$ and $c'(x, y)$ joining x, y using X_3, X_4 and X'_3, X_4 respectively (defined as before).

Let $V := U(X_1, X_2)$, $W := U(X_3, X_4)$ and $W' := U(X'_3, X_4)$. Then $V \subset W$, $V \subset W'$. Join $p(x)$ to V by a shortest geodesic γ in B and let $\tilde{\gamma}$, $\tilde{\gamma}'$ be the lifts of γ in X_3 and X'_3 respectively. Similarly join $p(x)$ to W, W' respectively by shortest geodesics $\gamma_{x,y}$ and $\gamma'_{x,y}$ and let $\tilde{\gamma}_{x,y}$ and $\tilde{\gamma}'_{x,y}$ be their lifts in X_3 and X'_3 respectively. Let $s_{x,y}$, $s'_{x,y}$ be the end points of $\tilde{\gamma}_{x,y}$, $\tilde{\gamma}'_{x,y}$ respectively, and let $b_{x,y}$, $b'_{x,y}$ be the end points of $\gamma_{x,y}$ and $\gamma'_{x,y}$.

Lemma 3.9. *With notation (in particular k, A) as above, there exists $D_{3.9}(= D_{3.9}(k, A))$ such that $d_B(b_{x,y}, b^{'}_{x,y})$ is bounded by $D_{3.9}$.*

Proof. Note that $V \subset W \cap W^{'}$, and that $V, W, W^{'}$ are all K-quasiconvex subsets of B. Therefore, by Lemma 1.24, concatenating $\gamma_{x,y}$ (resp. $\gamma^{'}_{x,y}$) with a geodesic joining $b_{x,y}$ (resp. $b^{'}_{x,y}$) to the terminal point of γ, we obtain $(3 + 2K)$-quasi-geodesics. These quasi-geodesics have the same end points as those of γ. Since B is δ-hyperbolic, therefore by Lemma 1.19 we can find $b, b^{'} \in \gamma$, such that $d_B(b_{x,y}, b) \leq D_3$, $d_B(b^{'}_{x,y}, b^{'}) \leq D_3$, where $D_3 := D_{1.19}(\delta, 3 + 2K)$. If $b \in [p(x), b^{'}] \subset \gamma$ then $b_{x,y} \in N_{2.D_3+\delta}(\gamma^{'}_{x,y})$. Otherwise, $b^{'} \in [p(x), b]$, so that $b^{'}_{x,y} \in N_{2.D_3+\delta}(\gamma_{x,y})$. So without loss of generality, we can assume that $b \in [p(x), b^{'}]$.

The end points of γ are in $U(X_3, X^{'}_3)$ which is a K-quasi-convex set in B. Hence by the bounded flaring condition (Corollary 1.12), we know that for all points $b_2 \in \gamma$, $d_{b_2}(X_3 \cap F_{b_2}, X^{'}_3 \cap F_{b_2}) \leq A.\mu(K)$. In particular, $d_b(X_3 \cap F_b, X^{'}_3 \cap F_b) \leq A.\mu(K)$. Similarly, $d_b(X_3 \cap F_b, X_4 \cap F_b) \leq A.\mu(D_3)$. Thus,

$$\begin{aligned} d_b(X^{'}_3 \cap F_b, X_4 \cap F_b) &\leq d_b(X_3 \cap F_b, X^{'}_3 \cap F_b) + d_b(X_3 \cap F_b, X_4 \cap F_b) \\ &\leq A.\mu(K) + A.\mu(D_3). \end{aligned}$$

We know that $[p(x), b^{'}] \subset N_{\delta+D_3}(\gamma^{'}_{x,y})$. Let $b^{'}_1 \in \gamma^{'}_{x,y}$ be such that $d_B(b, b^{'}_1) \leq \delta + D_3$. Then $d_{b^{'}_1}(X^{'}_3 \cap F_{b^{'}_1}, X_4 \cap F_{b^{'}_1}) \leq \mu(\delta + D_3).max\{d_b(X^{'}_3 \cap F_b, X_4 \cap F_b), 1\}$ and hence $d_{b^{'}_1}(X^{'}_3 \cap F_{b^{'}_1}, X_4 \cap F_{b^{'}_1}) \leq A.\mu(\delta + D_3)\{\mu(D_3) + \mu(K)\}$.

Denoting the right hand side of the preceding inequality by $D^{'}$, we have, by Lemma 2.17(2), $d_B(b^{'}_1, b^{'}_{x,y}) \leq D_{2.15}(k, D^{'})$. Since, $d_B(b_{x,y}, b^{'}_{x,y}) \leq d_B(b_{x,y}, b) + d_B(b, b^{'}_1) + d_B(b^{'}_1, b^{'}_{x,y})$, therefore

$$d_B(b_{x,y}, b^{'}_{x,y}) \leq D_3 + (\delta + D_3) + D_{2.17}(k, D^{'}) = \delta + 2D_3 + D_{2.17}(k, D^{'}).$$

Taking $D_{3.9} := \delta + 2D_3 + D_{2.17}(k, D^{'})$ completes the proof of the lemma. □

Lemma 3.10. *With k, A as above there exists $D_{3.10}(= D_{3.10}(k, A))$ such that the Hausdorff distance between $c(x, y)$ and $c^{'}(x, y)$ is bounded by $D_{3.10}$.*

Proof. **Step 1:** By Lemma 3.9 we have $d_B(b_{x,y}, b^{'}_{x,y}) \leq D_{3.9}(k, A)$. Hence, by δ-hyperbolicity of B, $Hd_B(\beta_{x,y}, \beta^{'}_{x,y}) \leq \delta + D_{3.9}(k, A)$. Since X_4 is a k-qi section, we have, by Remark 3.4(3),

$$Hd^{'}(\tilde{\beta}_{x,y}, \tilde{\beta}^{'}_{x,y}) \leq 2k.(\delta + D_{3.9}(k, A)).$$

Step 2: Similarly,

$$Hd^{'}([s_{x,y}, t_{x,y}], [s^{'}_{x,y}, t^{'}_{x,y}]) \leq A + 2k.D_{3.9}(k, A).$$

where $[s_{x,y}, t_{x,y}], [s^{'}_{x,y}, t^{'}_{x,y}]$ are the horizontal geodesic segments of $c(x, y)$ and $c^{'}(x, y)$ respectively, each of length at most A.

Step 3: Now we calculate the Hausdorff distance between $\tilde{\gamma}_{x,y}$ and $\tilde{\gamma}^{'}_{x,y}$. Let $\tilde{\gamma}^{''}_{x,y}$ be the lift of $\gamma_{x,y}$ in $X^{'}_3$. Then, as in Step 1, we have $Hd^{'}(\tilde{\gamma}^{'}_{x,y}, \tilde{\gamma}^{''}_{x,y}) \leq 2k.(\delta + D_{3.9}(k, A))$. Since γ joins two points $p(x)$ and $b_{x,y}$ of $U(X_3, X^{'}_3)$, it follows that $d_{b_2}(X_3 \cap F_{b_2}, X^{'}_3 \cap F_{b_2}) \leq A.\mu(K)$ for all points $b_2 \in \gamma$ (by Lemma 2.17(2) and the bounded flaring condition). Since there is a point $b \in \gamma$ such that $d_B(b, b_{x,y}) \leq$

$D_3 := D_{1.19}(\delta, 3+2K)$, we have the following using the boundedness of the flaring condition again:

$$\begin{aligned} d_{b_{x,y}}(X_3 \cap F_{b_2}, X_3^{'} \cap F_{b_2}) &\leq \mu(D_3).max\{d_b(X_3 \cap F_b, X_3^{'} \cap F_b), 1\} \\ &\leq A.\mu(D_3).\mu(K). \end{aligned}$$

Let $A_1 = A.\mu(D_3).\mu(K)$ and suppose $U_{A_1}(X_3, X_3^{'})$ is $K^{'} := D_{2.17}(c_2, A_1)$-quasi-convex. Note that $\gamma_{x,y}$ joins two points of $U_{A_1}(X_3, X_3^{'})$. Therefore, by Lemma 2.17 and the bounded flaring condition, we have $\forall b_1 \in \gamma_{x,y}$, $d_{b_1}(X_3 \cap F_{b_1}, X_3^{'} \cap F_{b_1}) \leq \mu(K^{'}).A_1$. Hence $Hd^{'}(\tilde{\gamma}_{x,y}, \tilde{\gamma}^{''}_{x,y}) \leq \mu(K^{'}).A_1$, and

$$\begin{aligned} Hd^{'}(\tilde{\gamma}_{x,y}, \tilde{\gamma}^{'}_{x,y}) &\leq Hd^{'}(\tilde{\gamma}_{x,y}, \tilde{\gamma}^{''}_{x,y}) + Hd^{'}(\tilde{\gamma}^{'}_{x,y}, \tilde{\gamma}^{''}_{x,y}) \\ &\leq \mu(K^{'}).A_1 + 2k.(\delta + D_{3.9}(k, A)). \end{aligned}$$

Finally, since

$$\begin{aligned} &Hd^{'}(c(x,y), c^{'}(x,y)) \\ &\leq \max\{Hd^{'}(\tilde{\beta}_{x,y}, \tilde{\beta}^{'}_{x,y}), Hd^{'}([s_{x,y}, t_{x,y}], [s^{'}_{x,y}, t^{'}_{x,y}]), Hd^{'}(\tilde{\gamma}_{x,y}, \tilde{\gamma}^{'}_{x,y})\}, \end{aligned}$$

the lemma follows, taking $D_{3.10} := \mu(K^{'}).A_1 + 2k.(\delta + D_{3.9}(k, A))$. ☐

Lemma 3.11. *With notation (in particular k, A) as above, there exists $D_{3.11}(= D_{3.11}(k, A))$ such that if $c(x,y), c(y,x)$ are defined using two k-qi sections X_3, X_4 where $k \leq c_4$, $x \in X_3$ and $y \in X_4$, then $Hd(c(x,y), c(y,x))$ is bounded by $D_{3.11}$.*

Proof. Let α be a geodesic in B joining $b_{x,y}$ and $b_{y,x}$. Since α joins two points of $U(X_3, X_4)$, we have $d_b(F_b \cap X_3, F_b \cap X_4) \leq \mu(K)A$ for all $b \in \alpha$, by Lemma 2.17(2) and the bounded flaring condition.

Since $U(X_3, X_4)$ is a K-quasi-convex set in B, $\gamma_{x,y} \cup \alpha$ is a $(3+2K)$-quasi-geodesic by Lemma 1.24. Hence, $Hd_B(\gamma_{x,y} \cup \alpha, [p(x), b_{y,x}]) \leq D_{1.19}(\delta, 3+2K)$, by Lemma 1.19. Similarly, $Hd_B(\gamma_{y,x} \cup \alpha, [p(y), b_{x,y}]) \leq D_{1.19}(\delta, 3+2K)$. It follows that $Hd^{'}(c(x,y), c(y,x)) \leq D_{3.11} := \mu(K)A + 2k(\delta + D_{1.19}(\delta, 3+2K))$. ☐

Corollary 3.12. *With notation (in particular k, A) as above, there exists $D_{3.12}(= D_{3.12}(k, A))$ such that the following holds.*
Let $x, y \in C(X_1, X_2)$ and $k \leq c_4$. Then the Hausdorff distance between any pair of curves joining x, y defined in the same way as $c(x,y)$ using k-qi sections passing through x, y, is at most $D_{3.12}$.

Proof. Choose $D_{3.12} := 2(D_{3.10} + D_{3.11})$. ☐

Lemma 3.13. *With notation (in particular k, A) as above with $k \leq c_4$, there exists $D_{3.13}(= D_{3.13}(k, A))$ such that the following holds.*
Suppose X_3, X_4, X_5 are k-qi sections in $C(X_1, X_2)$ such that $z \in X_5$, $y \in X_4 \subset C(X_1, X_5) \subset C(X_1, X_2)$ and $x \in X_3 \subset C(X_1, X_4) \subset C(X_1, X_2)$. Then the triangle formed by the curves $c(x,y), c(y,z), c(x,z)$, defined using the pairs X_3, X_4; X_4, X_5 and X_3, X_5 respectively, is $D_{3.13}-$slim.

Proof. We have $U(X_3, X_5) \subset U(X_4, X_5) \cap U(X_3, X_4)$ and we know, by Lemma 2.17, that all of these three sets are K-quasi-convex in B.

Case 1**:** *Suppose x, y are in the same horizontal space and $d_{p(x)}(x,y) \leq A$.*
Then $p(x) \in U(X_3, X_4)$. Since $\gamma_{x,z}$ ends in $U(X_3, X_5) \subset U(X_3, X_4)$, it joins two points of $U(X_3, X_4)$. Hence, by Lemma 2.17(2) and Corollary 1.12, we have for all $b^{'} \in \gamma_{x,z}$, $d_{b^{'}}(X_3 \cap F_{b^{'}}, X_4 \cap F_{b^{'}}) \leq A.\mu(K)$.

Now we show that $d_B(b_{x,z}, b_{y,z})$ is small. Recall that $b_{x,z} \in U(X_3, X_5) \subset U(X_4, X_5)$ and $b_{y,z} \in U(X_4, X_5)$. Thus $\gamma_{y,z} \cup [b_{y,z}, b_{x,z}]$ is a $(3+2K)$-quasi-geodesic in B, by Lemma 1.24. Hence, there is a point $b_2 \in \gamma_{x,z}$, such that $d_B(b_{y,z}, b_2) \leq D_{1.19}(\delta, 3+2K)$, by Lemma 1.19. Since $d_{b_{y,z}}(F_{b_{y,z}} \cap X_4, F_{b_{y,z}} \cap X_5) \leq A$, we have by bounded flaring, $d_{b_2}(F_{b_2} \cap X_4, F_{b_2} \cap X_5) \leq A.\mu(D_{1.19}(\delta, 3+2K))$. Therefore, $d_{b_2}(F_{b_2} \cap X_3, F_{b_2} \cap X_5) = d_{b_2}(F_{b_2} \cap X_3, F_{b_2} \cap X_4) + d_{b_2}(F_{b_2} \cap X_4, F_{b_2} \cap X_5) \leq A.\{\mu(K) + \mu(D_{1.19}(\delta, 3+2K)\}$. Now, by Lemma 2.17, we get

$$d_B(b_2, b_{x,z}) \leq D_{2.17}(k, A.\mu(K) + A.\mu(D_{1.19}(\delta, 3+2K)).$$

Hence

$$d_B(b_{x,z}, b_{y,z}) \leq D_{1.19}(\delta, 3+2K) + D_{2.17}(k, A.\mu(K) + A.\mu(D_{1.19}(\delta, 3))).$$

It follows by arguments similar to that of Lemma 3.10 that

$$Hd^{'}(c(x,z), c(y,z)) \leq A.\mu(K) + 2k.(\delta + d_B(b_{x,z}, b_{y,z})).$$

Replacing $d_B(b_{x,z}, b_{y,z})$ in the right hand expression with its upper bound obtained above, we get a constant $D^{'}_{3.13}$. The lemma follows, in this case, if we choose $D_{3.13} \geq D^{'}_{3.13}$.

Case 2: We consider the general case. For the rest of the proof, we shall assume that all the curves of the form $c(u,v)$ $(u, v \in X_3 \cup X_4 \cup X_5)$, are constructed using the sections X_3, X_4, X_5 only, unless otherwise specified. We first show that $Hd^{'}(c(x,z), \tilde{\gamma}_{x,y} \cup c(s_{x,y}, z))$ is bounded by a constant depending on k and A. Let $\bar{b}$ be a nearest point projection of $b_{x,y}$ on $U(X_3, X_5)$. By Lemma 1.24, $d_B(\bar{b}, b_{x,z}) \leq D_{1.24} = D_{1.24}(\delta, K)$. Let γ_2 be a geodesic joining $b_{x,y}$ to $\bar{b}$ and let $\widetilde{\gamma_2}$ be a lift of γ_2 in X_3. Note that $\gamma_{x,y} \cup \gamma_2$ is a $(3+2K)$-quasi geodesic in B. Thus the Hausdorff distance between $\gamma_{x,z}$ and $\gamma_{x,y} \cup \gamma_2$ is at most $\delta + D_{1.24} + D_{1.19}(\delta, 3+2K)$. Hence the Hausdorff distance between $\tilde{\gamma}_{x,y} \cup c(s_{x,y}, z)$ and $c(x,z)$, in $C(X_1, X_2)$, is at most $2k.\{\delta + D_{1.24} + D_{1.19}(\delta, 3+2K)\} + A = D_1$, say.

Again by case 1, we know that $Hd(c(s_{x,y}, z), c(t_{x,y}, z)) \leq D^{'}_{3.13}$. Thus $Hd(c(x,z), \tilde{\gamma}_{x,y} \cup [s_{x,y}, t_{x,y}] \cup c(t_{x,y}, z))$ is at most $A + D_1 + D^{'}_{3.13}$. It follows that if we define the curves $c(z, t_{x,y})$, $c(z,y)$ with respect to the sections X_4, X_5 by taking $\gamma_{z, t_{x,y}} = \gamma_{z,y}$, the triangle formed by the curves $c(z, t_{x,y})$, $c(z,y)$ and $\tilde{\beta}_{y, t_{x,y}}$ is $2k\delta$-slim.

Thus by Corollary 3.12, the triangle formed by the curves $\tilde{\beta}_{y, t_{x,y}}$, $c(t_{x,y}, z)$ and $c(y,z)$ is D_2-slim where $D_2 = 2k.\delta + 2.D_{3.12}$. Taking $D_{3.13} := A + D_1 + D^{'}_{3.13} + D_2$, the lemma follows. $\square$

Proof of Proposition 3.6

We verify that the set $\mathcal{S}$ of curves defined earlier in this section satisfies the properties of Corollary 1.29: Let $D = L$, $D_1 := 2c_2$. Fix any $D_2 \geq 1$ and $D_3 = D_{3.7}(c_2, A, D_2)$ and $D_4 = D_{3.13}(c_4, A) + 2.D_{3.12}(c_4, A)$.

Proof of properties 1 **and** 2**:** This is the content of Remark 3.4(2) and Lemma 3.7 respectively (also note the clarification in Remark 3.4(1)).

Proof of property 3**:** Suppose $x, y \in C(X_1, X_2)$. If $x^{'}, y^{'} \in c(x,y)$ then the segment of $c(x,y)$ between $x^{'}, y^{'}$, say $c(x,y)|_{[x^{'}, y^{'}]}$, is a possible candidate for the definition of $c(x^{'}, y^{'})$. Thus by Corollary 3.12, the Hausdorff distance of $c(x,y)|_{[x^{'}, y^{'}]}$ and $c(x^{'}, y^{'})$ is bounded by $D_{3.12}(c_2, A) \leq D_4$.

Proof of property 4**:** Let $x, y, z \in C(X_1, X_2)$. Then using Lemma 3.3 we may assume, without loss of generality, that x, y, z are contained in three c_4-qi sections X_3, X_4, X_5 respectively, where $X_4 \subseteq C(X_1, X_5)$, $X_3 \subseteq C(X_1, X_4)$. Now, the triangle formed by the curves $c(x, y)$, $c(y, z)$, $c(x, z)$ defined using these sections is $D_{3.13}(c_4, A)$-slim by Lemma 3.13. Hence, by Corollary 3.12, any triangle with vertices x, y, z formed by such curves is $\{D_{3.13}(c_4, A) + 2.D_{3.12}(c_4, A)\}$-slim. Let $\delta_{3.6} := \delta_{1.29}(D, D_1, D_2, D_3, D_4)$. It follows from Corollary 1.29, that $C(X_1, X_2)$ is $\delta_{3.6}$-hyperbolic.

By Remark 3.4(2), it follows that X_1, X_2 are the images of $2c_1-$quasi-isometric embeddings of B into the $\delta_{3.6}$-hyperbolic metric space $C(X_1, X_2)$. Thus, they are $K_{3.6} := D_{1.19}(\delta_{3.6}, 2c_1)$-quasiconvex in $C(X_1, X_2)$. This completes the proof of the first part of the proposition.

From the given conditions it follows by Lemma 2.17 that $U(X_1, X_2)$ is bounded. Hence, for any $x \in X_1$ and $y \in X_2$ the $K_{1.29}(D, D_1, D_2, D_3, D_4)$-quasi-geodesic $c(x, y)$ passes through the (uniformly) bounded set $p^{-1}(U(X_1, X_2)) \cap C(X_1, X_2)$. Since $C(X_1, X_2)$ has been proven to be hyperbolic, stability of quasi-geodesics (Lemma 1.19) completes the proof. $\square$

3.2. **Hyperbolicity of ladders: General case.**

Lemma 3.14. *There is a function $D_{3.14} : \mathbb{R}^+ \to \mathbb{R}^+$ such that the following holds. Suppose I, J are intervals in $\mathbb{R}$ and $\phi : I \to J$ is a k-quasi-isometric embedding. Let $x_1, x_2, x_3 \in I$, $x_1 \leq x_2 \leq x_3$, and suppose $\phi(x_1)$ belongs to the interval with end points $\phi(x_2)$, $\phi(x_3)$. Then $x_2 - x_1 \leq D_{3.14}(k)$.*

Proof. Without loss of generality, we may assume that $\phi(x_2) \leq \phi(x_1) \leq \phi(x_3)$. Let $x_4 = inf\{y \in [x_2, x_3] : \phi(y) \geq \phi(x_1)\}$.

If $x_2 = x_4$ then $\exists x^{'} \in [x_2, x_2 + 1] \cap [x_2, x_3]$ such that $\phi(x^{'}) \geq \phi(x_1)$. Now $x^{'} - x_2 \leq 1$ implies $|\phi(x^{'}) - \phi(x_2)| \leq 2k$, since ϕ is a k-quasi-isometric embedding. Therefore, $\phi(x_1) - \phi(x_2) \leq 2k$. Thus we have $x_2 - x_1 \leq 3k^2$.

If $x_2 < x_4$ we choose $x^{'} \in [x_2, x_4)$ and $x^{''} \in [x_4, x_3]$ such that $x_4 - x^{'} \leq 1$ and $x^{''} - x_4 \leq 1$ with $\phi(x^{''}) \geq \phi(x_1)$. Now $x^{''} - x^{'} \leq 2$ implies $|\phi(x^{'}) - \phi(x^{''})| \leq 3k$. Thus $\phi(x^{''}) - \phi(x_1) \leq 3k$, since $\phi(x^{'}) < \phi(x_1) \leq \phi(x^{''})$ by the choices of $x^{'}, x^{''}$. Hence $x_2 - x_1 \leq x^{''} - x_1 \leq 4k^2$. Therefore, in any case, we may choose $D_{3.14}(k) = 4k^2$. $\square$

Notation: For two qi-section X_1, X_2 in X and $D \geq 0$ we shall denote by $C_D(X_1, X_2)$ the D-neighborhood of the ladder $C(X_1, X_2)$ in X.

Lemma 3.15. *Given $f : \mathbb{R}^+ \to \mathbb{R}^+$, $k \geq 1, D \geq 2C_{3.3}(k)$, there exists $D^{'} \geq 1$ such that the following holds.*
Suppose $p : X \to B$ is an f-metric graph bundle and X_1, Y, X_2 are k-qi sections in X. Also suppose that Y is contained in the ladder $C(X_1, X_2)$. Then the D-neighborhood of each of the spaces Y, $C(X_1, Y), C(Y, X_2)$ is a connected subgraph of X and the intersection of the spaces $C_D(X_1, Y)$ and $C_D(Y, X_2)$ is contained in the $D^{'}$-neighborhood of Y in the path metric of both $C_D(X_1, Y)$ and $C_D(Y, X_2)$.

Proof. Since X_1, X_2, Y are k-qi sections and $D \geq 2C_{3.3}(k)$, it follows as was noted in Remark 3.4(1) that the D-neighborhood of each of the spaces $Y, C(X_1, Y), C(Y, X_2)$ is connected.

Now, let $y \in C_D(X_1, Y) \cap C_D(Y, X_2)$. Let us denote the path metric on $C_D(X_i, Y)$ induced from X by d_i and suppose $y_i \in C(X_i, Y)$ be such that $d_i(y, y_i) \leq D$, for

$i = 1, 2$. Then $d_B(p(y_1), p(y_2)) \leq 2D$. We need to prove the statements:
$\mathcal{P}_j$: *Any point of* $C_D(X_1, Y) \cap C_D(Y, X_2)$ *is contained in a* $D^{'}$*-neighborhood of* Y *in* $C_D(X_j, Y)$*, for* $j = 1, 2$.

Since the proofs of $\mathcal{P}_1, \mathcal{P}_2$ are similar, let us just prove $\mathcal{P}_2$. We know there exists a $k^{'} = C_{3.3}(k)$-qi section Y_2 say, through $y_2 \in C(X_2, Y)$ contained in $C(X_2, Y)$. Join y_2 to the point $y_1^{'} = Y_2 \cap F_{p(y_1)}$, by the lift of a geodesic in B joining $p(y_1)$ and $p(y_2)$. The length of this path is at most $4Dk^{'}$. Now, $d(y_1, y_1^{'}) \leq 2D + 4Dk^{'}$ and hence their horizontal distance is at most $f(2D + 4Dk^{'})$. Thus choosing $D^{'}$ to be $D + f(2D + 4Dk^{'})$, we are through. □

Suppose X_1, X_2 are two c_1-qi sections in X. Let us define the notation $c_{i+1} = C^i_{3.3}(c_1)$, $i \geq 1$, as in the proof of Proposition 3.6. Then we have the following.

Proposition 3.16. *For any* $L \geq 2c_6$*, and* $c_1 \geq 1$ *as above, there exists* $\delta_{3.16} = \delta_{3.16}(c_1, L)$ *such that* $C_L(X_1, X_2)$ *is a* $\delta_{3.16}$*- hyperbolic metric space with respect to the path metric induced from* X.

Proof. Let $A = K_1(M_{c_3^{'}} + K_1) + D_{3.14}(2g(c_3)) + f(2L + 4c_2.L)$. The idea of the proof is to break the ladder $C(X_1, X_2)$ into a finite number of subladders. Then by Proposition 3.6 and, if necessary, by a simple application Corollary 1.42 we show that these subladders are hyperbolic. Finally we apply Corollary 1.42 again to the ladder assembled out of subladders to finish the proof.

Step 1 : *Defining subladders.*
Fix a horizontal geodesic $\mathcal{I} = F_{b_0} \cap C(X_1, X_2)$. The two end points of $\mathcal{I}$ lie in X_1 and X_2. Choose a parametrization $\alpha : [0, l] \to \mathcal{I}$ by arc length so that $\alpha(0) \in X_1$ and $\alpha(l) \in X_2$. We shall inductively construct a finite sequence of integers $0 = s_0 < s_1 < \cdots < s_m = l$, and a sequence of c_2-qi sections $X_i^{'}$ contained in $C(X_1, X_2)$ such that $X_i^{'}$ passes through $\alpha(s_i)$ for each $i = 1, \cdots, m-1$. Let $X_0^{'} = X_1$. Suppose s_i has been obtained, $s_i < l$ and $X_i^{'}$ has been constructed. If $d_h(X_i^{'}, X_2) \leq A$ then define $s_{i+1} = l$, $X_{i+1}^{'} = X_2$ and the construction is over. Otherwise, consider the set

$$S_{i+1} = \{t \in [s_i, l] \cap \mathbb{N} : \exists \, \mathrm{a} \, c_2 - \mathrm{qi \ section} \, X^{'} \mathrm{through} \, \alpha(t) \, \mathrm{with} \, d_h(X^{'}, X_i^{'}) \leq A\}$$

Let $u_{i+1} = \max S_{i+1}$. If $\exists t \in S_{i+1}$ such that there is a c_2-qi section $X^{'}$ inside $C(X_1, X_2)$ through $\alpha(t)$ with $d_h(X^{'}, X_i^{'}) = A$, define $s_{i+1} = t$ and $X_{i+1}^{'} = X^{'}$. Otherwise define $s_{i+1} = min\{l, u_{i+1} + 1\}$ and let $X_{i+1}^{'}$ be any c_2-qi section inside $C(X_1, X_2)$ through $\alpha(s_{i+1})$. The construction of these sections stops at the m-th step if $d_h(X_{m-1}^{'}, X_2) \leq A$, so that we must have $X_m^{'} = X_2$ and $s_m = l$. It follows from the above construction of the sections $X_i^{'}$ that for each i, $1 \leq i \leq m-1$, we have $d_h(X_{i-1}^{'}, X_i^{'}) \geq A$ and in case $d_h(X_i^{'}, X_{i+1}^{'}) > A$, there is a section $X_i^{''}$ through a point $\alpha(t_i)$, $t_i \in [s_i, s_{i+1}]$ with $d_h(X_j^{'}, X_i^{''}) \leq A$, $j = i, i+1$.

Step 2 : *Subladders form a decomposition of* $C(X_1, X_2)$.
In this step, we will show that $C(X_1, X_2) = \cup_{i=0}^m C(X_i^{'}, X_{i+1}^{'})$ and that $C(X_{i-1}^{'}, X_i^{'}) \cap C(X_i^{'}, X_{i+1}^{'}) = X_i^{'}$.

Note that the first assertion follows from the second and the construction in Step 1, and for the second assertion, it is enough to show the following:
Claim: $X_{i+1}^{'} \subseteq C(X_i^{'}, X_2)$, for all i, $1 \leq i \leq m-2$.

Consider the triples of points $(X_1 \cap F_b, X_i^{'} \cap F_b, X_{i+1}^{'} \cap F_b)$, $b \in \mathcal{V}(B)$. They are contained in the geodesic $F_b \cap C(X_1, X_2)$. For $b = b_0$ we know, by the construction in Step 1, that $X_i^{'} \cap F_b \in [X_1 \cap F_b, X_{i+1}^{'} \cap F_b]$.

We now argue by contradiction. Suppose $X_{i+1}^{'} \not\subseteq C(X_i^{'}, X_2)$. Then for some point $b^{'} \in \mathcal{V}(B)$, we must have $X_{i+1}^{'} \cap F_{b'} \in [X_1 \cap F_{b'}, X_i^{'} \cap F_{b'}]$. Therefore there exist $b_1, b_2 \in \mathcal{V}(B)$ with $d(b_1, b_2) = 1$, such that $X_i^{'} \cap F_{b_1} \in [X_1 \cap F_{b_1}, X_{i+1}^{'} \cap F_{b_1}]$ but $X_{i+1}^{'} \cap F_{b_2} \in [X_1 \cap F_{b_2}, X_i^{'} \cap F_{b_2}]$. We know that $X_i^{'}, X_{i+1}^{'}$ are c_2-quasi-isometric sections, and X_1 is a c_1-quasi-isometric section. Hence $d(X_i^{'} \cap F_{b_1}, X_i^{'} \cap F_{b_2}) \leq 2c_2$, $d(X_{i+1}^{'} \cap F_{b_1}, X_{i+1}^{'} \cap F_{b_2}) \leq 2c_2$ and $d(X_1 \cap F_{b_1}, X_1 \cap F_{b_2}) \leq 2c_1 \leq 2c_2$.

By Lemma 3.3, the definition of c_3 (at the top of the proof of this proposition) and Lemma 1.10, we have a $g(2c_3)$-quasi-isometric embedding $[X_1 \cap F_{b_1}, X_{i+1}^{'} \cap F_{b_1}] \to [X_1 \cap F_{b_2}, X_2 \cap F_{b_2}]$ which sends each of the points $X_j^{'} \cap F_{b_1}$ to $X_j^{'} \cap F_{b_2}$, $j = i, i+1$ and $X_1 \cap F_{b_1}$ to $X_1 \cap F_{b_2}$. By Lemma 3.14 we get

$$d_{b_1}(X_i^{'} \cap F_{b_1}, X_{i+1}^{'} \cap F_{b_1}) \leq D_{3.14}(g(2c_3)).$$

By the choice of the constant A, and the definition of $X_i^{'}$'s this gives rise to a contradiction, completing the proof of Step 2.

Step 3 : *Subladders are uniformly hyperbolic.*
Next we show that there are constants δ_1, k_1 and D such that each $C_L(X_i^{'}, X_{i+1}^{'})$ is δ_1-hyperbolic and $X_i^{'}, X_{i+1}^{'}$ are k_1-quasi-convex in $C_L(X_i^{'}, X_{i+1}^{'})$ for each i, $0 \leq i \leq m-1$. Also we shall show that the sets $X_i^{'}, X_{i+1}^{'}$ are mutually D-cobounded in $C_L(X_i^{'}, X_{i+1}^{'})$, for $i \neq m-1$.

Since $X_i^{'}, X_{i+1}^{'}$ are c_2-qi sections in X, it follows that they are the images of $2c_2-$quasi-isometric embeddings in $C_L(X_i^{'}, X_{i+1}^{'})$ (Remark 3.4(2)). Hence, they will be $D_{1.19}(\delta_1, 2c_2)$-quasiconvex in $C_L(X_i^{'}, X_{i+1}^{'})$ provided we can show that $C_L(X_i^{'}, X_{i+1}^{'})$ is δ_1-hyperbolic.

If $d_h(X_i^{'}, X_{i+1}^{'}) \leq A$ then, by Proposition 3.6, each $C_L(X_i^{'}, X_{i+1}^{'})$ is $\delta_{3.6}(c_2, A, L)$-hyperbolic; moreover, in this case, unless $i = m-1$, we have $d_h(X_i^{'}, X_{i+1}^{'}) = A$ and $X_i^{'}, X_{i+1}^{'}$ are then mutually $D_{3.6}(c_2, A, D)$-cobounded.

Suppose $d_h(X_i^{'}, X_{i+1}^{'}) > A$. Recall that $X_j^{'}$ passes through $\alpha(s_j)$, $j = i, i+1$. In this case, we can find $t_i \in [s_i, s_{i+1}]$ such that there is a c_2-qi section $X_i^{''}$ in $C(X_1, X_2)$, passing through $\alpha(t_i)$, so that $d_h(X_j^{'}, X_i^{''}) \leq A$, $j = i, i+1$. Now, as in the proof of Lemma 3.3, we project points of $X_i^{''}$ into the horizontal geodesics of $C(X_i^{'}, X_{i+1}^{'})$ and get a c_3-qi section $Y_i^{'}$ through $\alpha(t_i)$. Note that we still have $d_h(X_j^{'}, Y_i^{'}) \leq A$ for $j = i, i+1$. By Proposition 3.6, $C_L(X_i^{'}, Y_i^{'})$, and $C_L(X_{i+1}^{'}, Y_i^{'})$ are both $\delta_{3.6}(c_3, A, L)$-hyperbolic. Also we see that $C_L(X_i^{'}, Y_i^{'}) \cap C_L(X_{i+1}^{'}, Y_i^{'})$ contains a $2c_3$-neighborhood of $Y_i^{'}$ which is connected. Since $Y_i^{'}$ is a c_3-quasi-isometric image of B in X, hence it is a $2c_3$-quasi-isometric image in both $C_L(X_i^{'}, Y_i^{'})$ and $C_L(X_{i+1}^{'}, Y_i^{'})$.

Now, we apply Lemma 3.15 followed by Corollary 1.42. Here the total space is $C_L(X_i^{'}, X_{i+1}^{'})$ and we have just two subspaces: $C_L(X_i^{'}, Y_i^{'})$ and $C_L(X_{i+1}^{'}, Y_i^{'})$. Also their intersection contains a $2c_3$-neighborhood of $Y_i^{'}$, denoted by Y_i, say.

Thus, $C_L(X^{'}_i, X^{'}_{i+1})$ is $\delta_{1.42}(\delta_{3.6}(c_3, A, L), D^{'}_{3.15}(f, c_3, L), 2c_3, 1)$-hyperbolic. Define

$$\delta_1 := max\{\delta_{3.6}(c_2, A, L), \delta_{1.42}(\delta_{3.6}(c_3, A, L), D^{'}_{3.15}(f, c_3, L), 2c_3, 1)\}.$$

We next show that the quasi-convex sets $X^{'}_i, X^{'}_{i+1}$ are mutually cobounded in $C_L(X^{'}_i, X^{'}_{i+1})$ in this case too. Due to the symmetry of the situation, it suffices to show that the nearest point projection of $X^{'}_{i+1}$ on $X^{'}_i$ is uniformly bounded.

Since the sets $U(X^{'}_j, Y^{'}_i)$, $j = i, i+1$ are K-quasiconvex in B, the lift Y_{ij} (say) of $U(X^{'}_j, Y^{'}_i)$ in $Y^{'}_i$ is a $C_1 := (2Kc_3 + D_{1.19}(\delta_1, 2c_3))$-quasi-convex set in $C_L(X^{'}_i, X^{'}_{i+1})$. Let $R = R_{1.26}(\delta_1, C_1) + L_{1.21}(3 + 2C_1, 3 + 2C_1, \delta_1)$. Now we have two cases:

(a) If Y_{ij}, $j = i, i+1$ are not R-separated then there are two points in $X^{'}_i$ and $X^{'}_{i+1}$ respectively which are at a distance of at most $A^{'}_1 := (2A + R)$ from each other. Then, it follows by Lemma 3.7 that $d_h(X^{'}_i, X^{'}_{i+1}) \leq A_1 := f(2A^{'}_1 c_2 + A^{'}_1)$, by the definition of a metric graph bundle. Hence, by Proposition 3.6, $X^{'}_i, X^{'}_{i+1}$ are $D_{3.6}(c_2, A_1)$-cobounded.

(b) Else suppose Y_{ij}, $j = i, i+1$ are R-separated. In this case we note the following:

1. Any geodesic in $C_L(X^{'}_i, X^{'}_{i+1})$ joining two points of Y_{ij}, $j = 1, 2$, passes close to the end points of the coarsely unique geodesic joining these sets. Thus we will be done if we can show that any geodesic joining $X^{'}_j$, $j = 1, 2$ passes close to the end points of this coarsely unique geodesic.
2. Any nearest point projection of a point in $X^{'}_j$ on $Y^{'}_i$ is uniformly close to the set Y_{ij} for $j = i, i = 1$.
3. Let $x \in X^{'}_{i+1}$ and let $y \in X^{'}_i$ be a nearest point projection of x on $X^{'}_i$. Let x_1 be a nearest point projection of x on Y_{ii+1} and let y_1 be a nearest point projection of y on Y_{ii}. Then the union of the three geodesic segments $[x, x_1], [x_1, y_1], [y, y_1]$ is a uniform quasi-geodesic. Hence the Hausdorff distance between this quasi-geodesic and a geodesic $[x, y]$ is uniformly bounded. But $[x_1, y_1]$ passes close to the end points of the coarsely unique geodesic joining the sets Y_{ij}, $j = 1, 2$.

It follows that any geodesic joining $X^{'}_j$, $j = 1, 2$ passes close to the end points of this coarsely unique geodesic and step 3 follows.

Step 4 : *The final step:*

Finally we use Lemma 3.15 in conjunction with Corollary 1.42. Here the total space is $C_L(X_1, X_2)$, and the sequence of subspaces are $C_L(X^{'}_i, X^{'}_{i+1})$, $i = 0, 1, \ldots, m-1$. We check to see that the hypotheses of Corollary 1.42 are satisfied:

(1) Each of the subspaces $X^{'}_i$ is δ_1-hyperbolic by step 3;

(2) by choice of the constant $A > f(2L + 4c_2.L)$ (see Lemma 3.7) we know that only the consecutive ones intersect nontrivially;

(3) for $i = 2, \ldots, m$, the intersection of two consecutive subspaces $C_L(X^{'}_j, X^{'}_{j+1})$, $j = i-1, i$, contains the $2c_2$-neighborhood, say Y_i, of $X^{'}_i$ which is connected (Lemma 3.7); and also the intersection is contained in the $D^{'}_{3.15}(f, c_3, L)$-neighborhood of Y_i in the spaces $X^{'}_j$, j=1,2

(4) Y_i are $2c_2$-quasi-isometrically embedded since $X^{'}_i$ are c_2-quasi-isometrically embedded in X

(5) the sets X_i', X_{i+1}' are uniformly cobounded in $C_L(X_i', X_{i+1}')$ for $i = 1, 2, \ldots, m-2$ as proved in Step 3.

The proposition follows. $\square$

4. The Combination Theorem

In this section we prove the main theorem of our paper which says that *a metric (graph) bundle satisfying the properties mentioned at the beginning of Section 3 has hyperbolic total space* (see paragraph entitled "Notations and conventions" at the beginning of Section 3).

As in Section 3, we replace a metric bundle by an approximating metric graph bundle. Thus we shall work exclusively with metric graph bundles and apply the results of the last section. Here is an outline of the main steps of the proof.
For any pair of points $x, y \in X$ we choose a ladder containing x, y and choose a geodesic $c(x, y)$ joining x, y in a connected neighborhood of the ladder. Thus we have a family of curves and we show that the family satisfies the conditions of Corollary 1.29. Proofs of conditions 1 and 2 follow from the results of the last section. Proofs of conditions 3 and 4 follow from Proposition 4.2 below, which says that large neighborhoods of 'tripod bundles' are hyperbolic. This in turn follows from Proposition 3.16 and Corollary 1.42.

Definition 4.1. *For three qi sections X_1, X_2, X_3 in a metric graph bundle X over B a* tripod bundle *determined by these qi sections, denoted $C(X_1, X_2, X_3)$, is defined to be the union of the ladders $C(X_1, X_2)$, $C(X_2, X_3)$, $C(X_3, X_1)$.*

The convention that we adopted after Definition 3.1 applies here as well; namely, since the Hausdorff distance between any two tripod bundles determined by three qi sections is uniformly bounded we denote by $C(X_1, X_2, X_3)$ any tripod bundle determined by the qi sections X_1, X_2, X_3. Also for any qi sections X_1, X_2, X_3 in X and $D \geq 0$ we denote by $C_D(X_1, X_2, X_3)$ the D-neighborhood of the tripod bundle $C(X_1, X_2, X_3)$ in X.

4.1. **Proof of Main Theorem.**

Suppose X over B is an (f, K)-metric graph bundle. We continue with the notation and conventions fixed at the beginning of section 3. In particular,
1) X is either a metric graph bundle satisfying a flaring condition or one obtained as an approximating metric graph bundle of a metric bundle satisfying a flaring condition;
2) the base space and horizontal spaces are uniformly hyperbolic;
3) the barycenter maps of the horizontal spaces are uniformly surjective.

Now, suppose X_1,X_2 and X_3 are three c_1-qi sections in X. We shall show that for large $D \geq 0$, $C_D(X_1, X_2, X_3)$ is hyperbolic. For this, we first show that taking a nearest point projection of $X_3 \cap F_b$ onto the horizontal geodesic $C(X_1, X_2) \cap F_b$ (for all $b \in B$) we get a qi section X_4. (See figure below.) Then we have a genuine 'tripod bundle' $C(X_1, X_2) \cup C(X_3, X_4)$, such that $C_D(X_1, X_2, X_3)$ is quasi-isometric to an L-neighborhood of $C(X_1, X_2) \cup C(X_3, X_4)$, where L depends on D and the bundle. The quasi-isometry is provided by projecting any point z of $C_L(X_1, X_2) \cup C_L(X_3, X_4)$ onto a nearest point in $C_D(X_1, X_2, X_3)$ lying in the same horizontal fiber as z (Here, the nearest point-projection is taken in the metric on the horizontal fiber to which z belongs.) Hyperbolicity of the space $C_L(X_1, X_2) \cup C_L(X_3, X_4)$, and quasi-convexity of $C(X_1, X_2)$ in this space follow essentially from Proposition 3.16 and Corollary 1.42.



Tripod

Proposition 4.2. *Given* $c_1 \geq 1$*, there exists* $L_0, \delta_{4.2}, K_{4.2} \geq 0$ *such that the following holds.*
Let X_1, X_2, X_3 *be* c_1*-qi sections and* $L \geq L_0$*. Then* $C_L(X_1, X_2, X_3)$ *is* $\delta_{4.2}(= \delta_{4.2}(c_1, L))$*-hyperbolic with the induced path metric from* X *and each of* $C_L(X_i, X_j)$*,* $i \neq j$ *is* $K_{4.2}(= K_{4.2}(c_1, L))$*-quasi-convex in* $C_L(X_1, X_2, X_3)$*.*

For ease of exposition, we break the proof up into several lemmas, many of which will be minor modifications of results we have shown already.

For $b_1, b_2 \in B$ with $d(b_1, b_2) = 1$, we have a $g(2c_1)$-quasi-isometry $F_{b_1} \to F_{b_2}$ by Lemma 1.10, which sends $X_i \cap F_{b_1}$ to $X_i \cap F_{b_2}$ for $i = 1, 2, 3$. Therefore, by Lemma 1.27, choosing a nearest point projection of $X_3 \cap F_b$ onto the horizontal geodesic $[X_1 \cap F_b, X_2 \cap F_b]$, for all $b \in B$, we get a $c_1^{'}$-qi section of B in X where $c_1^{'} := 2c_1 + D_{1.27}(\delta^{'}, g(2c_1))$. Let us call this section X_4. Let $c_{i+1}^{'} := C_{3.3}^{i}(c_1^{'})$, $i \geq 1$.

Now we have the following analog of Lemma 3.15.

Lemma 4.3. *For all* $L \geq 2c_2^{'}$*, there exists* $D_{4.3}(= D_{4.3}(L))$ *such that the intersection* $C_L(X_1, X_2) \cap C_L(X_3, X_4)$ *is contained in the* $D_{4.3}$*-neighborhood of* X_4*.*

Proof. The proof is an exact copy of that of the proof of Lemma 3.15, hence we omit the details. The only extra observation needed is that the curve $[X_3 \cap F_b, X_4 \cap F_b] \cup [X_4 \cap F_b, X_i \cap F_b]$, $i = 1, 2$ is a $(3, 0)$-quasi-geodesic in F_b. $\square$

Lemma 4.4. *For all* c_1 *as above and* $L \geq 2c_6^{'}$*, there exist* $D_{4.4}(= D_{4.4}(c_1, L))$ *and* $K_{4.4}(= K_{4.4}(c_1, L))$ *such that the space* $C_L(X_1, X_2) \cup C_L(X_3, X_4)$ *is* $D_{4.4}$*-hyperbolic and* $C(X_1, X_2)$ *is* $K_{4.4}$*-quasi-convex in this space.*

Proof. The first part of the lemma follows as an application of Proposition 3.16 and Corollary 1.42 (the proof is a replica of Step 3 of the proof of Proposition 3.6 which shows that large girth subladders are hyperbolic). For completeness we briefly check the conditions of Corollary 1.42.

(1) Here we have only two subgraphs $C_L(X_1, X_2)$ and $C_L(X_3, X_4)$ which are hyperbolic by Proposition 3.6.
(2) Condition (2) is trivially satisfied.
(3) The intersection $C_L(X_1, X_2) \cap C_L(X_3, X_4)$ contains the $c_2^{'}$-neighborhood say, Y, of X_4 which is connected and the rest follows from Lemma 4.3 above.
(4) Since X_4 is $2c_1^{'}$-quasi-isometrically embedded in $C_L(X_1, X_2) \cup C_L(X_3, X_4)$, Y is also quasi-isometrically embedded.
(5) Condition 5 is trivially satisfied.

For the second part of the lemma we note that any geodesic joining two points of $C(X_1, X_2)$ in $C_L(X_1, X_2) \cup C_L(X_3, X_4)$ and which leaves $C_L(X_1, X_2)$ must join two points in a (uniformly bounded) neighborhood of X_4, by Lemma 4.3. Since X_4 is the image of a quasi-isometric embedding of B in the hyperbolic space $C_L(X_1, X_2) \cup C_L(X_3, X_4)$ it is quasi-convex also. The lemma follows. ☐

Clearly for all $b \in \mathcal{V}(B)$, $C(X_1, X_2, X_3) \cap F_b$ is $\delta^{'}$-quasi-convex in F_b. Define a map $\Pi : Z = C_L(X_1, X_2) \cup C_L(X_3, X_4) \to X$ by sending any point $x \in Z \cap F_b$ to a nearest point in $C(X_1, X_2, X_3) \cap F_b$ (in the d_b-metric).

Lemma 4.5. *Given $c_1 \geq 1$ there exists $D_{4.5}(= D_{4.5}(c_1))$ such that the map Π is $D_{4.5}-$Lipschitz.*

Proof. We need to check that for any two adjacent vertices in the domain of Π, the image vertices are at a uniformly bounded distance. This breaks up into two cases.

When the vertices are in the same horizontal space F_b then since $C(X_1, X_2, X_3) \cap F_b$ is (uniformly) quasiconvex in F_b, and since nearest point projections onto quasiconvex sets in hyperbolic metric spaces are coarsely Lipschitz (cf. Lemma 3.2 of [Mit98b]) the claim follows.

When the vertices are not in same horizontal space then the same argument as in Lemma 1.27 [Mit98b, Bow07] shows that nearest-point projections and quasi-isometries almost commute. The rest of the proof is a replica of Theorem 3.5 [Mit98a]. ☐

Remark 4.6. *In fact, it follows easily that Π restricted to $C(X_1, X_2)$ is a qi-embedding into any sufficiently large neighborhood of $C(X_1, X_2, X_3)$ where the latter is equipped with a path metric induced from X.*

Lemma 4.7. *Given $c_1 \geq 1$ and $L \geq 0$ as above there exists $D_{4.7}(= D_{4.7}(c_1, L))$ such that the following holds.*
For all $x \in C_L(X_1, X_2) \cup C_L(X_3, X_4)$ the horizontal distance between x and $\Pi(x)$ is at most $D_{4.7}$.

Proof. This follows from the fact that in any $\delta'-$ hyperbolic metric space (=F_b in our case) the Hausdorff distance between a triangle with vertices x, y, z and the tripod $[x, w] \cup [y, z]$ (where $w \in [y, z]$ is a nearest point projection of x onto $[y, z]$) is bounded by δ'. ☐

Proof of Proposition 4.2:
Set $L_0 = 2c_6^{'} + D_{4.5}(c_1)$; by assumption $L \geq L_0$. Let $L_1 = L + \delta'$.

First for every pair of points in $x, y \in C(X_1, X_2, X_3)$ we choose a geodesic in $C_{L_1}(X_1, X_2) \cup C_{L_1}(X_3, X_4)$ and project it into $C(X_1, X_2, X_3)$ by Π. This defines a path in $C(X_1, X_2, X_3)$ say $c(x, y)$ joining x, y. Note that by Lemma 4.7 the paths $c(x, y)$ are *uniform quasi-geodesics in* $C_{L_1}(X_1, X_2) \cup C_{L_1}(X_3, X_4)$. Now we need to check the conditions of Corollary 1.29.

Here the whole space is $C_L(X_1, X_2, X_3)$ and the discrete set is the set of vertices contained in $C(X_1, X_2, X_3)$. Next we note the following:

1. Conditions (1), (2) of Corollary 1.29 follow from Lemma 4.5.
2. Conditions (3), (4) of Corollary 1.29 follow from Lemma 4.7, since the space $C_{L_1}(X_1, X_2) \cup C_{L_1}(X_3, X_4)$ has been proved to be hyperbolic.

This shows that $C_L(X_1, X_2, X_3)$ is hyperbolic. From Lemmas 4.4, and 4.7 it follows that $C(X_1, X_2)$ is the image of the quasi-convex set $C(X_1, X_2) \subset C_{L_1}(X_1, X_2) \cup C_{L_1}(X_3, X_4)$ under the quasi-isometric embedding Π (cf. Remark 4.6). Hence it is quasi-convex in $C_L(X_1, X_2, X_3)$ and thus so is $C_L(X_1, X_2)$.

This completes the proof. □

Let $\Pi_1 : C_L(X_1, X_2, X_3) \to C(X_1, X_2)$ be defined by the same prescription as that of Π, namely for all $b \in \mathcal{V}(B)$ and $x \in C_L(X_1, X_2, X_3) \cap F_b$, $\Pi_1(x)$ is defined to be a nearest point projection of x onto the horizontal geodesic $C(X_1, X_2) \cap F_b$. We have the following corollary of Proposition 4.2.

Corollary 4.8. *Given* $c_1 \geq 1$ *and* $L \geq L_0$ *(cf. Proposition 4.2) there exists* $D_{4.8}(= D_{4.8}(c_1, L))$ *such that if*
1) $x, y \in C_L(X_1, X_2)$
2) γ_1 *is a geodesic in* $C_L(X_1, X_2, X_3)$ *joining* x, y.
3) γ_2 *is a geodesic in* $C_L(X_1, X_2)$ *joining* x, y
Then the Hausdorff distance $Hd(\gamma_1, \gamma_2) \leq D_{4.8}$.

Proof. For simplicity of notation, denote the distance function of $C_L(X_1, X_2, X_3)$ also by d. For all $z \in \gamma_1$, there exists $z_1 \in C_L(X_1, X_2)$ such that $d(z, z_1) \leq K_{4.2}(c_1, L)$. Hence there exists $z_2 \in C(X_1, X_2)$ such that $d(z, z_2) \leq L + K_{4.2}(c_1, L)$. Therefore, there exists $D_1 \geq 0$ depending on c_1, L such that $d(z, \Pi_1(z)) \leq D_1$. Thus $\Pi_1(\gamma_1)$ is a $(1, 2D_1)-$quasi-geodesic in $C_L(X_1, X_2, X_3)$.

We claim that $\Pi_1(\gamma_1)$ is also a (uniform) quasi-geodesic in $C_L(X_1, X_2)$. To show this, let d_1 denote the path metric on $C_L(X_1, X_2)$ induced from X. Then Π_1 is Lipschitz by the same argument used to show that Π is Lipschitz in Lemma 4.5. Suppose $x_1, x_2 \in \gamma_1$ and let $y_i = \Pi_1(x_i)$, $i = 1, 2$. Then $d_1(y_1, y_2) \geq d(y_1, y_2) \geq d(x_1, x_2) - 2D_1$. The claim follows.

Hence $\Pi_1(\gamma_1)$ and γ_2 are (uniform) quasi-geodesics in $C_L(X_1, X_2)$ joining x, y. Since $C_L(X_1, X_2)$ is hyperbolic the Hausdorff distance between $\Pi_1(\gamma_1)$ and γ_2 is uniformly bounded. Similarly, since $C_L(X_1, X_2, X_3)$ is hyperbolic, it follows that the Hausdorff distance between $\Pi_1(\gamma_1)$ and γ_1 is uniformly bounded. The Corollary follows. □

Corollary 4.9. *Given* $c_1 \geq 1$ *and* $L \geq 0$ *there exists* $D_{4.9}(= D_{4.9}(c_1, L))$ *such that the following holds.*
Suppose $X_i, X_i^{'}$, $i = 1, 2$ *are* c_1*-qi sections and* $x_i \in X_i \cap X_i^{'}$, $i = 1, 2$. *Then the Hausdorff distance between the geodesics joining* x_1, x_2 *in the subspaces* $C_L(X_1, X_2)$ *and* $C_L(X_1^{'}, X_2^{'})$ *is at most* $D_{4.9}(c_1, L)$.

Proof. This follows from Proposition 4.2 and Corollary 4.8 applied successively to the tripod bundles $C_L(X_1, X_1^{'}, X_2)$ and $C_L(X_1^{'}, X_2, X_2^{'})$. □

Now, we have all the ingredients we need to prove our main combination theorem.

Theorem 4.10. *Suppose $p : X \to B$ is a metric bundle (resp. metric graph bundle) such that*
(1) *B is a δ-hyperbolic metric space.*
(2) *Each of the fibers F_b, $b \in B$ (resp. $b \in \mathcal{V}(B)$) is a $\delta^{'}$-hyperbolic metric space with respect to the path metric induced from X.*
(3) *The barycenter maps $\partial^3 F_b \to F_b$, $b \in B$ (resp. $b \in \mathcal{V}(B)$) are (uniformly) coarsely surjective.*
(4) *A flaring condition is satisfied.*
Then X is a hyperbolic metric space.

Proof. Let c_1 be as above (the constant associated with the qi section), and set $L = L_0$ as in Proposition 4.2. Now, we define a set of curves joining each pair of points $x, y \in X$ and check that they satisfy the properties of Corollary 1.29 to complete the proof.

Definition of the curves: For each pair of points x, y in X, choose, once and for all, two c_1-qi sections X_1, X_2 passing through x and y respectively. Now define $c(x, y)$ to be a geodesic in $C_L(X_1, X_2)$ joining x, y.

- **Proof of property 1:** This follows by taking $D_1 = 1$.
- **Proof of property 2:** This follows from Proposition 3.6 and Lemma 3.7. One just needs to recall that the curves constructed in the proof of Proposition 3.6 are uniform quasi-geodesics.
- **Proof of property 3:** This follows from Proposition 4.2, and Corollary 4.8.
- **Proof of property 4:** Given $x, y, z \in X$ choose three c_1-qi sections X_3, X_4, X_5 containing $x, y, z \in X$ respectively and define the curves $c^{'}(x, y)$, $c^{'}(x, z)$ and $c^{'}(y, z)$ using these sections in the same way as the curves $c(x, y)$ are defined. It follows from Corollary 4.8 that the triangle formed by $c^{'}(x, y)$, $c^{'}(x, z)$ and $c^{'}(y, z)$ is $(\delta_{4.2}(c_1, L) + 2D_{4.8}(c_1, L))$-slim. Corollary 4.9 now gives property 4.

□

Remark 4.11. 1. Let us now unravel the hypotheses. Note that for proving that tripod bundles are hyperbolic what we crucially used are
a) ladders are hyperbolic (Proposition 3.16)
b) horizontal spaces F_b are uniformly hyperbolic.

In order to prove that the ladders are hyperbolic what we needed are
c) the base space B is hyperbolic
d) the conclusion of Lemma 2.17 holds.

Finally to construct ladders we needed to prove the existence of qi sections (Proposition 2.10).

2. Note that the conditions of Theorem 4.10 are inherited by induced metric graph bundles over quasi-isometrically embedded subsets of B. Hence the induced bundles over quasi-isometrically embedded subsets of B are also hyperbolic.

5. Consequences and Applications

A number of consequences of Theorem 4.10 are collected together in this section.

5.1. **Sections, Retracts and Cannon-Thurston maps.** We shall say that an exact sequence of finitely generated groups $1 \to K \to G \to Q \to 1$ satisfies *bounded flaring* if the associated metric graph bundle (cf. Example 1.5) of Cayley graphs does. An immediate consequence of Theorem 4.10 coupled with the existence of qi-sections from Theorem 2.11 is the following *converse* to Mosher's Theorem 2.11.

Theorem 5.1. *Suppose that the short exact sequence of finitely generated groups*

$$1 \to K \to G \to Q \to 1.$$

satisfies a flaring condition such that K, Q *are word hyperbolic and* K *is non-elementary. Then* G *is hyperbolic.*

Theorem 2.11 was generalized by Pal [Pal10] as follows.

Theorem 5.2. (Pal [Pal10]**)** *Suppose we have a short exact sequence of finitely generated groups*

$$1 \to K \to G \overset{p}{\to} Q \to 1,$$

with K *strongly hyperbolic relative to a subgroup* K_1 *such that* G *preserves cusps, i.e. for all* $g \in G$ *there exists* $k \in K$ *with* $gK_1g^{-1} = kK_1k^{-1}$. *Then there exists a* $(k, \epsilon)-$*quasi-isometric section* $s\colon Q \to G$ *for some constants* $k \geq 1, \epsilon \geq 0$.

Next, suppose we have a short exact sequence of pairs of finitely generated groups

$$1 \to (K, K_1) \to (G, N_G(K_1)) \overset{p}{\to} (Q, Q_1) \to 1$$

with K *(strongly) hyperbolic relative to the cusp subgroup* K_1. *If* G *preserves cusps, then* $Q_1 = Q$ *and there is a quasi-isometric section* $s\colon Q \to N_G(K_1)$ *satisfying*

$$\frac{1}{R} d_Q(q, q') - \epsilon \leq d_{N_G(K_1)}(s(q), s(q')) \leq R d_Q(q, q') + \epsilon$$

where $q, q' \in Q$ *and* $R \geq 1, \epsilon \geq 0$ *are constants. Further, if* G *is weakly hyperbolic relative to* K_1*, then* Q *is hyperbolic.*

The setup of Theorem 5.2 naturally gives a metric graph bundle $P : X \to Q$ of spaces, where Q is the quotient group and fibers are the coned off spaces $\widehat{K}$ obtained by electrocuting copies of K_1 in K.

We shall now use a version of Theorem 3.5. Theorem 3.5 is proven in [Mit98a] in the context of an exact sequence of finitely generated groups $1 \to N \to G \to Q \to 1$, with N hyperbolic; but all that the proof requires is the existence of qi sections (which follows in the context of groups by the qi section Theorem 2.11 of Mosher).

As in [Mit98a], the existence of a qi-section through each point of X guarantees, via Theorem 3.5, the existence of a continuous extension to the boundary (also called a Cannon-Thurston map [CT07] [CT85]) of the map $i_b : F_b \to X$ provided X is hyperbolic. The proof is identical to that in [Mit98a] and we omit it here, referring the reader to [Mit98a] for details. Combining this fact with Theorem 4.10 we have the following.

Theorem 5.3. *Suppose* $p : X \to B$ *is a metric (graph) bundle with the following properties:*

(1) B *is a* δ*-hyperbolic metric space.*

(2) *Each of the fibers* F_b, $b \in B$ *is a* δ'*-hyperbolic metric space with respect to the induced path metric from* X.

(3) *The barycenter maps* $\partial^3 F_b \to F_b$, $b \in B$ $(b \in \mathcal{V}(B))$ *are uniformly coarsely*

surjective.
(4) *The metric (graph) bundle satisfies a flaring condition.*
Then the inclusion $i_b : F_b \to X$ *extends continuously to a map* $\hat{i} : \widehat{F_b} \to \widehat{X}$ *between the Gromov compactifications.*

5.2. **Hyperbolicity of base and flaring.** In our main combination theorem 4.10 flaring was a sufficient condition. In this subsection and the next we investigate its necessity. This issue is closely linked with hyperbolicity of the base space B. We study it with special attention to hyperbolic and relatively hyperbolic groups as in Theorems 2.11 and Theorem 5.2.

A Theorem of Papasoglu (cf. [Pap95], Lemma 3.8 of [Pap05]) states the following.

Theorem 5.4. [Pap95, Pap05] *Let* G *be a finitely generated group and let* Γ *be the Cayley graph of* G *with respect to to a finite generating set. If there is an* ϵ *such that geodesic bigons in* Γ *are* ϵ*-thin then* G *is hyperbolic.*
Similarly, let X *be a geodesic metric space such that for every* K *there exists* C *such that* K*-quasigeodesic bigons are* C*-thin, then* X *is hyperbolic.*
In fact there is some (universal) constant $C > 0$ *such that if* G *is finitely generated and non-hyperbolic, then* $\forall R > 0$ *there is some* $R' > R$ *and a* (C, C)*-quasi-isometric embedding of a Euclidean circle of radius* R' *in* Γ*.*

We now look at short exact sequences of finitely generated groups.

Proposition 5.5. *Consider a short exact sequence of finitely generated groups*

$$1 \to K \to G \to Q \to 1.$$

such that K *is non-elementary word hyperbolic but* Q *is not hyperbolic. Then the short exact sequence cannot satisfy a flaring condition.*

Proof. By Theorem 5.4, Q contains (C, C) qi embeddings of Euclidean circles of arbitrarily large radius. Now, given any l, A_0 construct
a) a (C, C) qi embedding τ_l of a Euclidean circle σ of circumference $> 4l$ in Q
b) two qi sections s_1, s_2 of Q into G by Theorem 2.11 such that $d_h(s_1 \circ \tau_l(\sigma), s_2 \circ \tau_l(\sigma)) > A_0$.

Let $q \in \sigma$ be such that the length $d_q(s_1 \circ \tau_l(q), s_2 \circ \tau_l(q))$ in the fiber F_q over q is maximal. Let the two arcs of length l in τ_l starting at q (in opposite directions) end at q_1, q_2. Let $\overline{q_1 q q_2}$ denote the union of these arcs. Then the two quasigeodesics $s_1 \circ \tau_l(\overline{q_1 q q_2}), s_2 \circ \tau_l(\overline{q_1 q q_2})$ violate flaring as the fiber length achieves a maximum at the midpoint q. □

We next turn to the relatively hyperbolic situation described in Theorem 5.2 with Q non-hyperbolic, i.e. we assume that K is (strongly) hyperbolic relative to K_1. We have an analog of Proposition 5.5 in this situation too. The proof is the same as that of the above Proposition. The existence of qi sections in this case, follows from Theorem 5.2.

Lemma 5.6. *Suppose we have a short exact sequence of finitely generated groups*

$$1 \to (K, K_1) \to (G, N_G(K_1)) \xrightarrow{p} (Q, Q_1) \to 1$$

such that K *strongly hyperbolic relative to the cusp subgroup* K_1 *and* G *preserves cusps, but* Q *is not hyperbolic. Let* $P : X \to Q$ *be the associated metric graph bundle of spaces, where* Q *is the quotient group and fibers* F_q *are the coned off spaces* $\widehat{K}$ *obtained by electrocuting copies of* K_1 *in* K*. Then* X *does not satisfy flaring.*

The rest of this subsection is devoted to proving the following.

Proposition 5.7. *Suppose we have a short exact sequence of finitely generated groups*

$$1 \to (K, K_1) \to (G, N_G(K_1)) \xrightarrow{p} (Q, Q_1) \to 1$$

with K (strongly) hyperbolic relative to the cusp subgroup K_1 such that G preserves cusps. Suppose further that G is (strongly) hyperbolic relative to $N_G(K_1)$. Then Q is hyperbolic.

Proof. We shall argue by contradiction. Suppose Q is not hyperbolic.

Let X be a Cayley graph of G with respect to a finite generating set S containing a finite generating set of K (and, for good measure, a finite generating set of K_1). Let B be the Cayley graph of Q with respect to $p(S)$. Then the quotient map $G \to Q$ gives rise to a metric graph bundle $p : X \to B$ as before. This metric graph bundle admits uniform qi sections through each point of X by Theorem 5.2. Also B is not a hyperbolic metric space. By Theorem 5.4, there exists $C > 0$ such that for all $r > 0$ we can construct a (C, C)-qi embedding τ_r of a Euclidean circle σ_r of radius bigger than r in B.

Claim: Given $k > 0$ there exists D such that for any k-qi section $s : \mathcal{V}(B) \to X$ of the metric bundle $p : X \to B$, $s \circ \tau_r(\sigma_r)$ is contained in a D-neighborhood of a coset of $N_G(K_1)$.

Proof of claim: Let $\tau = s \circ \tau_r$. Then τ is a $k_1 := (kC + k)$-quasi-isometric embedding of σ_r in $s(B)$. Let u, v be a pair of antipodal points of the circle and $a = \tau(u), b = \tau(v)$. Let σ_r^1, σ_r^2 be the two arcs of σ_r joining u, v. Then $\tau(\sigma_r^1), \tau(\sigma_r^2)$ are k_1-quasigeodesics joining a, b.

Let d_r denote the intrinsic path metric on σ_r. Since $\sigma_r^1 \cup \sigma_r^2$ is a quasigeodesic bigon, it follows that for all $C_1 \geq 0$, there exists $C_2 \geq 0$ such that for all $r > 0$ and $x, y \in \sigma_r$, $d_r(x, \{a, b\}) \geq C_2$ and $d_r(y, \{a, b\}) \geq C_2$ implies that $d_X(x, y) \geq C_1$. Hence $\tau(\sigma_r^1) \cup \tau(\sigma_r^2)$ is a 'thick' quasigeodesic bigon, i.e. except for initial and final subsegments of length $k_1 C_2$, $\tau(\sigma_r^1)$ and $\tau(\sigma_r^2)$ are separated from each other by at least $\frac{C_1}{k_1}$.

Since G is strongly hyperbolic relative to $N_G(K_1)$, thick geodesic bigons lie in a bounded neighborhood of a coset of $N_G(K_1)$ [Far98]. The claim follows. □

We continue with the proof of the proposition. For any $k-$qi section $s : \mathcal{V}(B) \to X$, we shall refer to $s \circ \tau_r(\sigma_r) = \tau(\sigma_r)$ as a qi section of the circle σ_r. Let Y_1, Y_2 be two k-qi sections of a large Euclidean circle σ_r in B, such that Y_1 and Y_2 lie $D-$close to two *distinct* cosets of $N_G(K_1)$ (with $D = D(k)$ as in the Claim above). Let $W(Y_1, Y_2)$ be the union $\cup_{q \in \tau_r(\sigma_r)} \lambda_q$, where λ_q is a horizontal geodesic in F_q joining $Y_1 \cap F_q$ to $Y_2 \cap F_q$. Suppose $b, b^{'}$ are images (under τ_r) of antipodal points on σ_r. As in the proof of Lemma 3.3, we know that there exists $k_1 (= k_1(k))$ such that for each point z of λ_b there exists a k_1-quasi-isometric section of σ_r in $W(Y_1, Y_2)$; any such qi section is $D_1 (= D_1(k1))-$close to a coset of $N_G(K_1)$ by the Claim above.

Since Y_1, Y_2 are close to distinct cosets of $N_G(K_1)$, we can find (as in Step 1 of Proposition 3.16) two k_1-qi sections $Y_1^{'}$, $Y_2^{'}$ of σ_r passing through $z_1, z_2 \in \lambda_b$ with $d(z_1, z_2) = 1$ such that $Y_1^{'}$, $Y_2^{'}$ are

a) D_1-close to two distinct cosets of $N_G(K_1)$,

b) both contained in $W(Y_1, Y_2)$.

Suppose $Y_1^{'}, Y_2^{'}$ intersect $\lambda_{b'}$ in $z_1^{'}$ and $z_2^{'}$ respectively. If $d(z_1^{'}, z_2^{'})$ is large, in the same way as before, we can construct two $k_2(= k_2(k_1))-$qi sections Y_3, Y_4 of σ_r contained in $W(Y_1^{'}, Y_2^{'})$ such that
a) Y_3, Y_4 are $D_2(= D_2(D_1))-$ close to two distinct cosets of $N_G(K_1)$
b) $d(Y_3 \cap \lambda_{b'}, Y_4 \cap \lambda_{b'}) = 1$.

Thus we have two k_2-qi sections of long subarcs of σ_r that start and end close by in X but lie close to distinct cosets of $N_G(K_1)$. Since the r can be chosen to be arbitrarily large, this violates strong relative hyperbolicity of G with respect to the cosets of $N_G(K_1)$, proving the proposition. □

5.3. **Necessity of Flaring.** In this subsection we prove that flaring is a necessary condition for hyperbolicity of a metric (graph) bundle:

Proposition 5.8. *Let $P : X \to B$ be a metric (graph) bundle such that*
1. *X is δ-hyperbolic*
2. *There exist δ_0 such that each of the fibers F_z, $z \in B(\mathcal{V}(B))$ is δ_0-hyperbolic equipped with the path metric induced from X.*
Then the metric bundle satisfies a flaring condition.

In particular, any exact sequence of finitely generated groups $1 \to N \to G \to Q \to 1$ with N, G hyperbolic satisfies a flaring condition.

The proof will occupy the entire subsection. Suppose $\gamma : [-L, L] \to B$ is a geodesic and α, β are two K_1-qi lifts of γ. As in the construction of ladders, we define Y to be the union of horizontal geodesics $[\alpha(t), \beta(t)] \subset F_{\gamma(t)}$, $t \in [-L, L]$, and refer to it as the *ladder* formed by α and β. Let $\eta : [0, M] \to F_{\gamma(0)}$ be the geodesic $Y \cap F_{\gamma(0)}$.

A crucial ingredient is the following lemma which is a specialization to our context of the fact that geodesics in a hyperbolic space diverge exponentially. (See Proposition 2.4 and the proof of Theorem 4.11 in [Mit97]).

Lemma 5.9. *Given $K_1 \geq 1, D \geq 0$ there exist $b = b(K_1, D) > 1$, $A = A(K_1, D) > 0$ and $C = C(K_1, D) > 0$ such that the following holds:*

If $d(\alpha(0), \beta(0)) \leq D$ and there exists $T \in [0, L]$ with $d(\alpha(T), \beta(T)) \geq C$ then any path joining $\alpha(T + t)$ to $\beta(T + t)$ and lying outside the union of the $\frac{T+t-1}{2K_1}$-balls around $\alpha(0), \beta(0)$ has length greater than Ab^t for all $t \geq 0$ such that $T + t \in [0, L]$. In particular, the horizontal distance between $\alpha(T + t)$ and $\beta(T + t)$ is greater than Ab^t for all $t \geq 0$ such that $T + t \in [0, L]$.

Now, we use Lemma 5.9 to show that the ladder Y flares in at least one direction of γ. We start the proof by showing this in two special cases. A general ladder is then broken into subladders of the special types by qi lifts of γ as in Step 1 of the proof of Proposition 3.16. (Recall that we get exactly two types of subladders in this way. This motivates us to consider the two types of special ladders here.) We point out that

(1) the first type of ladder is of uniformly small (but not too small) girth;

(2) the second type of ladder is not necessarily of small girth but any qi lift of γ divides it into two subladders of small girth.

The proof of flaring for all ladders follows from this.

We shall need the following lemma also.

Lemma 5.10. *Given $d_1, d_2, \delta \geq 0$ and $k \geq 1$ there are constants $C = C(d_1, d_2, k, \delta)$ and $D = D(k, \delta)$ such that the following holds:*

Let X be a δ-hyperbolic metric space and let $\alpha_1, \alpha_2 : [-L, L] \to X$ be k-quasi-geodesics. Let $[a, b] \subset [-L, L]$ and suppose $d_1 = d(\alpha_1(a), \alpha_2(a))$ and $d_2 = d(\alpha_1(b), \alpha_2(b))$. If $[t - C, t + C] \subset [a, b]$ for some $t \in [a, b]$ then $d(\alpha_1(t), \alpha_2(t)) \leq D$.

This follows easily from stability of quasi-geodesics and slimness of triangles in X. (See Lemma 1.15, Chapter $III.H$, [BH99] for instance).

Remark 5.11. *1. We shall assume L to be sufficiently large for the following arguments to go through.*

2. As in the proof of Lemma 3.3 we can show that through each point of a ladder Y formed by K_1-qi lifts of a geodesic γ in B there is a $C_{3.3}(K_1)$-qi lift of γ contained in Y.

Flaring of ladders in special cases:

Let $D = D_{5.10}(K_1, \delta)$ and $D_1^{'} = C_{5.9}(K_1, D)$. Since the horizontal spaces in X are uniformly properly embedded in X there is a D_1 such that for all $v \in B$ and $x, y \in F_v$ if $d_v(x, y) \geq D_1$ then $d(x, y) \geq D_1^{'}$. Let $K_{i+1} = C_{3.3}^{i}(K_1)$.

Lemma 5.12. Ladders of type 1: *For K_1, D as above and $M \geq D_1$, there exists $n_1 = n_1(K_1, M)$ such that $max\{d_{\gamma(-t)}(\alpha(-t), \beta(-t)), d_{\gamma(t)}(\alpha(t), \beta(t))\} \geq 4M$ for all $t \geq n_1$.*

Proof. Suppose that $d_{\gamma(0)}(\alpha(0), \beta(0)) = M \geq D_1$. Let $D_2 = C_{5.9}(K_1, M)$ and let $C_1 := 1 + 2.C_{5.10}(M, D_2, K_1, \delta)$. If $d(\alpha(C_1), \beta(C_1)) \geq D_2$ then for all $t \geq 0$ the length of the horizontal geodesic joining $\alpha(C_1 + t)$ to $\beta(C_1 + t)$ is greater than or equal to $A_1.b_1^t$ for some $A_1 = A_{5.9}(M, K_1)$, $b_1 = b_{5.9}(M, K_1)$. Choose $t_1 > 0$ such that for all $t \geq t_1$, $A_1.b_1^{t_1} \geq 4.M$.

Else, suppose $d(\alpha(C_1), \beta(C_1)) < D_2$. In this case, by Lemma 5.10, $d(\alpha(\frac{C_1 - 1}{2}), \beta(\frac{C_1 - 1}{2})) \leq D$. By the choice of the constants D, D_2 we can again apply Lemma 5.9 so that for all $t \geq 0$ the length of a horizontal geodesic $Y \cap F_{\gamma(-t)}$ is greater than or equal to $A_2.b_2^t$, where the constants A_2, b_2 depend on K_1 and D. Choose $t_2 > 0$ such that for all $t \geq t_2$, $A_2.b_2^t \geq 4.M$. Now let $n_1 = max\{C_1 + t_1, t_2\}$. Thus we have $max\{d_{\gamma(-t)}(\alpha(-t), \beta(-t)), d_{\gamma(t)}(\alpha(t), \beta(t))\} \geq 4M$ for all $t \geq n_1 = n_1(M, K_1)$. $\square$

Lemma 5.13. Ladders of type 2: *Suppose $l > 0$ and that for any $s \in [0, M - 1]$ there is a K_2- qi lift α_1 of γ in Y through $\eta(s)$ such that $d(\alpha(t), \alpha_1(t)) \leq l$ for some $t \in [-L, L]$. There are $n_2 = n_2(K_1, l)$ and $D_4 = D_4(K_1, l)$ such that for all $t \geq n_2$ we have*

$$max\{d_{\gamma(t)}(\alpha(t), \beta(t)), d_{\gamma(-t)}(\alpha(-t), \beta(-t))\} \geq 4.M \text{ if } M - 1 \geq D_4.$$

Proof. Let $C_3 = C_{5.9}(K_3, l)$, $A_3 = A_{5.9}(K_3, l), b_3 = b_{5.9}(K_3, l)$. Let $m_0 := min\{m \in \mathbb{N} : A_3.b_3^m \geq D_{3.14}(g(2K_3))\}$, where g refers to the function defined in Lemma 1.10. It follows easily from the bounded flaring condition that there is a constant $D_4^{'}$ such that the following is true:

Suppose we have two K_3-qi lifts $\alpha^{'}, \alpha^{''} : [-L, L] \to X$ of the geodesic $\gamma : [-L, L] \to B$ with $d_{\gamma(0)}(\alpha^{'}(0), \alpha^{''}(0)) \geq D_4^{'}$ then $d(\alpha^{'}(t), \alpha^{''}(t)) \geq D_{3.14}(g(2K_3))$ for all $t \in [0, m_0]$.

Let $D_4 = max\{D_4^{'}, C_3\}$ and let $M - 1 = N.D_4 + r$ where $N \in \mathbb{N}$ and $0 \leq r < D_4$. Now construct a K_2-qi section β_1 in the ladder Y such that $d_{\gamma(0)}(\beta(0), \beta_1(0)) = r + 1$ and $d(\alpha(t_0), \beta_1(t_0)) \leq l$ for some $t_0 \in [-L, L]$. Without loss of generality we assume that $t_0 \in [-L, 0]$. We now break the subladder of Y, formed by α and

β_1, by K_3-qi lifts $\alpha_0 = \alpha, \alpha_1, \cdots, \alpha_N = \beta_1$ of γ such that $d_{\gamma(0)}(\alpha_i(0), \alpha_{i+1}(0)) = D_4$. We have $d(\alpha_i(t_0), \alpha_{i+1}(t_0)) \leq l$. Thus by the choice of the constant D_4, $d_{\gamma(t)}(\alpha_i(t), \alpha_{i+1}(t)) \geq max\{D_{3.14}(g(2K_3)), A_3.b_3^t\}$ and (as in Step 2 of the proof of Proposition 3.16) $\cup[\alpha_i(t), \alpha_{i+1}(t)]$ is a partition of the horizontal geodesic segment $[\alpha(t), \beta_1(t)] \subset Y \cap F_{\gamma(t)}$, for all $t \in [0, m_0]$. Therefore, we can choose $n_2 = n_2(K_1, l)$ such that for all $t \geq n_2$,

$$max\{d_{\gamma(t)}(\alpha(t), \beta(t)), d_{\gamma(-t)}(\alpha(-t), \beta(-t))\} \geq 4.M \text{ if } M - 1 \geq D_4 = D_4(K_1, l).$$

□

Flaring of general ladders: In the general case, first of all, we break the ladder Y into subladders of special types as described above (see Figure below where horizontal and vertical directions have been interchanged for aesthetic reasons).

Let us assume that Y is bounded by $K-$ qi lifts α, β of a geodesic $\gamma : [-L, L] \rightarrow X$. Let $\eta : [0, M] \rightarrow F_{\gamma(0)}$ be the geodesic $Y \cap F_{\gamma(0)}$. Let $K_i = C^i_{3.3}(K)$, and $l = D_{3.14}(g(2.K_2))$. Let $M_K := max\{D_1(K_1), D_4(K_1, l)\}$, and $n_K := max\{n_1(K_1, D_1(K_1)), n_2(K_1, l)\}$ where the functions D_1, D_4, n_1, n_2 are as in the proof of the flaring for the special ladders.

Claim: If $M \geq M_K$ then we have

$$max\{d_{\gamma(-n_K)}(\alpha(-n_K), \beta(-n_K)), d_{\gamma(n_K)}(\alpha(n_K), \beta(n_K))\} \geq 2.d_{\gamma(0)}(\alpha(0), \beta(0)).$$

To show this we inductively construct K_1-qi sections $\alpha_0 = \alpha, \alpha_1, \cdots, \alpha_i = \beta$ in Y to decompose it into subladders of the two types we mentioned above. This is done as in Step 1 of the proof of Proposition 3.16. Nevertheless we include a sketch of the argument for completeness.

Let $M \geq M_K$. By Remark 5.11 (2), we can construct a K_1-qi section α_1 through $\eta(M_K)$. Now, suppose $\alpha_1, \cdots, \alpha_j$ has been constructed through the points $\eta(s_1), \cdots, \eta(s_j)$ respectively. If $d_{\gamma(0)}(\alpha_j(0), \beta(0)) \leq M_K$ define $\alpha_{j+1} = \beta$. Otherwise, if there is a K_1-qi section through $\eta(M_K + s_j)$ in the ladder formed by α_j and β define it to be α_{j+1}. If neither of that happens then consider the following set:

$$\mathcal{T}_j = \{t \geq s_j + M_K : \exists \text{ a } K_1\text{-qi section through } \eta(t) \text{ entering the ladder formed by } \alpha_j \text{ and } \alpha\}$$

Let $t_j = \sup \mathcal{T}_j$ be the supremum of this set. Define α_{j+1} to be a K_1-qi section through $s_{j+1} := t_j + 1$, in the ladder formed by α_j and β that does not enter the ladder formed by α_j and α.



Flaring subladders

For each j, α_j and α_{j+1} form a special ladder and hence it must flare. Thus η can be expressed as the disjoint union of subsegments that flare to the left and the union of the subsegments that flare to the right respectively. The total length of one of these types must be at least half of the length of η. The claim follows. □

The first statement of Proposition 5.8 follows immediately. The last statement follows from Example 1.5 and the first part of this Proposition. □

5.4. **An Example.** Let $(Teich(S), d_T)$ be the Teichmuller space of a closed surface S equipped with the Teichmuller metric d_T. Teichmuller space can also be equipped with an electric metric d_e by electrocuting the cusps (see [Far98] for details on electric geometry and the introduction to this paper for a quick summary and notation). Note (as per work of Masur-Minsky [MM99], see also [Mj09]) that $(Teich(S), d_e)$ is quasi-isometric to the curve complex $CC(S)$. Let $E : (Teich(S), d_T) \rightarrow (Teich(S), d_e)$ be the identity map from the Teichmuller space of S equipped with the Teichmuller metric d_T to the Teichmuller space of S equipped with the electric metric d_e.

We shall need the following Theorem due to Hamenstadt [Ham10] which used an idea of Mosher [Mos03] in its proof.

Theorem 5.14. Hamenstadt [Ham10]**:** *For every $L > 1$ there exists $D > 0$ such that the following holds.*
Let $f : \mathbb{R} \rightarrow (Teich(S), d_T)$ be a Teichmuller L-quasigeodesic such that $E \circ f : \mathbb{R} \rightarrow (Teich(S), d_E)$ is also an L-quasigeodesic. Then for all $a, b \in \mathbb{R}$ there is a Teichmuller geodesic η_{ab} joining $f(a), f(b) \in Teich(S)$ such that the Hausdorff distance $d_H(f([a,b]), \eta_{ab}) \leq D$.

We are now in a position to prove a rather general combination Proposition for metric bundles over quasiconvex subsets of $CC(S)$.

Proposition 5.15. *Let (K, d_K) be a hyperbolic metric space satisfying the following,*
There exists $C > 0$ such that for any two points $u, v \in K$, there exists a bi-infinite C-quasigeodesic $\gamma \subset K$ with $d_K(u, \gamma) \leq C$ and $d_K(v, \gamma) \leq C$.
Let $j : K \rightarrow (Teich(S), d_T)$ be a quasi-isometric embedding such that $E \circ j : K \rightarrow (Teich(S), d_e)$ is also a quasi-isometric embedding. Let $U(S, K)$ be the pull-back bundle under j of the universal curve over $Teich(S)$ equipped with the natural path metric. Then $\widetilde{U(S,K)}$ is a hyperbolic metric space. Here $\widetilde{U(S,K)}$ denotes the metric bundle obtained by taking fiberwise universal covers.

Proof. Clearly, $\widetilde{U(S,K)}$ is a metric bundle over K (since the universal curve over $Teich(S)$ is topologically a product $S \times Teich(S)$). Hence, by Theorem 4.10 it suffices to prove flaring. Let S_x denote the fiber of $U(S, K)$ over $x \in i(K)$.

Let $[a, b]$ be a geodesic segment of sufficiently large length in K. By the hypothesis on K, there exists a bi-infinite geodesic passing within a bounded neighborhood of $[a, b]$. Hence without loss of generality, we may assume that a, b lie on a bi-infinite geodesic in K.

By Theorem 5.14 we may assume that $j(a), j(b)$ lie in a uniformly bounded neighborhood of a fat Teichmuller geodesic η_{ab} whose end-points in $\partial Teich(S)$ are two singular foliations $\mathcal{F}_+, \mathcal{F}_-$. Let d_s be the singular Euclidean metric on S induced by the pair of singular foliations $\mathcal{F}_+, \mathcal{F}_-$.

The rest of the argument follows an argument of Mosher [Mos97]. Let x be some point on the fat Teichmuller geodesic η_{ab} obtained in the previous paragraph. Given any geodesic segment $\lambda \subset \widetilde{S_x}$ of length $l(\lambda)$, there are two projections λ_+ and λ_- onto (the universal covers of) $\mathcal{F}_+, \mathcal{F}_-$ in $\widetilde{S_x}$. At least one of these projections is of length at least $\frac{l(\lambda)}{2}$. If u, v are two points on either side of x such that $d_T(u,x) \geq m$ and $d_T(v,x) \geq m$, then the length of λ in at least one of $\widetilde{S_u}$ and $\widetilde{S_v}$ is greater than $e^m \frac{l(\lambda)}{2}$.

Since all the surfaces with piecewise Euclidean metric involved in the above argument can be chosen to have uniformly bounded diameter, their universal covers are all uniformly quasi-isometric to a fixed Cayley graph of $\pi_1(S)$. Flaring follows. $\square$

Remark 5.16. The same proof goes through for S^h- a finite area surface with cusps, *provided we equip the cusps with a zero metric.* (This is the electric metric on cusps in the terminology of [Far98].) Flaring in this situation is proved in Section 4.4 of [MR08].

From Remark 5.16 we obtain directly the following consequence (see [FM02] for definitions).

Consider a surface S^h with punctures. Let $K = \pi_1(S^h)$ and let $\mathcal{K} = \{K_1, \cdots, K_p\}$ be the collection of peripheral subgroups. The pure mapping class group is the subgroup of the mapping class group that preserves individual punctures. Let Q be a convex cocompact subgroup of the pure mapping class group. We state the Proposition below for a surface with a single puncture for convenience where the pure mapping class group is the mapping class group.

Proposition 5.17. *Let $K = \pi_1(S^h)$ be the fundamental group of a surface with a single puncture and K_1 be its peripheral subgroup. Let Q be a convex cocompact subgroup of the mapping class group of S^h. Let*

$$1 \to (K, K_1) \to (G, N_G(K_1)) \xrightarrow{p} (Q, Q_1) \to 1$$

be the induced short exact sequence of (pairs of) groups. Then G is (strongly) hyperbolic relative to $N_G(K_1)$.

Conversely, if G is (strongly) hyperbolic relative to $N_G(K_1)$, then Q is convex-cocompact.

Proof. Suppose that Q is convex cocompact. Then Q is hyperbolic by [FM02], [KL08]. Also $Q = Q_1$ by Theorem 5.2. Let $\mathcal{E}(G, K_1)$ denote the electric space obtained from (the Cayley graph of) G after coning off translates of (the Cayley graph of) K_1. Note that $\mathcal{E}(G, K_1)$ is a metric graph bundle quasi-isometric to a $\widehat{K}(= \mathcal{E}(K, K_1))$-bundle over Q where $\widehat{K}(= \mathcal{E}(K, K_1))$ denotes K with copies of K_1 coned off. The flaring condition for this bundle and hence weak relative hyperbolicity of the pair (G, K_1) follow from Remark 5.16.

Let $\mathcal{Q}$ denote the collection of translates of (Cayley graphs of) $Q(= N_G(K_1)/K_1)$ in $\mathcal{E}(G, K_1)$, where each copy of Q in $\mathcal{E}(G, K_1)$ is a copy of the electric space $\mathcal{E}(N_G(K_1), K_1)$ obtained by coning off K_1 in translates of (Cayley graphs of) $N_G(K_1)$.

To prove that G is (strongly) hyperbolic relative to $N_G(K_1)$ it suffices to prove that $\mathcal{E}(G, K_1)$ is (strongly) hyperbolic relative to $\mathcal{Q}$, as K is already strongly hyperbolic relative to K_1 by [Far98]. This in turn would follow [Mj11] from (uniform) mutual coboundedness of pairs of elements in $\mathcal{Q}$. Note also that each $Q_i \in \mathcal{Q}$ is quasi-isometrically embedded and hence a quasiconvex subset of $\mathcal{E}(G, K_1)$. Any two such Q_1, Q_2's define a ladder $C(Q_1, Q_2)$ by regarding Q_1, Q_2 as qi sections of the metric graph bundle $\mathcal{E}(G, K_1)$ over Q. Each $C(Q_1, Q_2)$ is hyperbolic by Proposition 3.16. Hence, the ladder $C(Q_1, Q_2)$ also satisfies flaring by Proposition 5.8.

We now argue by contradiction. Let d_h denote the horizontal distance in $\mathcal{E}(G, K_1)$. Suppose that elements of the collection $\mathcal{Q}$ do not satisfy (uniform) mutual coboundedness. Then there exists $D_0 > 0$ such that for any $l > 0$, there exists a pair $Q_1, Q_2 \in \mathcal{Q}$ and a geodesic segment $r : [0, l] \to Q$ such that $d_h(s_1 \circ r(t), s_2 \circ r(t)) \leq D_0$ for all $t \in [0, l]$, where $s_i : Q \to \mathcal{E}(G, K_1)$ are quasi-isometric embeddings defining the sections Q_1, Q_2. Since the number of elements in K of length at most D_0 is bounded it follows that for sufficiently large l, there exists $q \in Q, q \neq 1$ and $h \in (K \setminus K_1), h \neq 1$ such that $s(q), h$ commute. This is impossible for Q convex cocompact, proving the forward direction of the Proposition.

We now prove the converse direction. Hyperbolicity of Q follows from Proposition 5.7. To prove convex cocompactness, it is enough to show by [FM02] that *some* orbit of the action of Q on $(Teich(S), d_T)$ is quasiconvex.

Since G is (strongly) hyperbolic relative to $N_G(K_1)$, it follows from Lemma 1.39 that $\mathcal{E}(G, K_1)$ is strongly hyperbolic relative to the collection $\mathcal{Q}$ of translates of (Cayley graphs of) $Q(= N_G(K_1)/K_1)$ in $\mathcal{E}(G, K_1)$ as defined above. Since Q is hyperbolic, it follows (cf. [Bow97] Section 7) that $\mathcal{E}(G, K_1)$ is hyperbolic. Thus, $\mathcal{E}(G, K_1)$ is a hyperbolic metric graph bundle over Q. Hence, from Proposition 5.8, the bundle $\mathcal{E}(G, K_1)$ over Q satisfies flaring. The logarithm of the stretch factor guaranteed by flaring gives a lower bound on the Teichmuller distance.

The remainder of the argument is an exact replica of the proof of Theorem 1.2 of [FM02] (Section 5.2 in particular), which proves the analogous statement for surfaces without punctures. We do not reproduce the argument here but point out that the only place in the proof where explicit use is made of closedness of S is Theorem 5.5 of [FM02], which, in turn is taken from [Mos03]. A straightforward generalization of this fact to the punctured surface case is given in [Pal11]. $\square$

Remark 5.18. *It is worth noting that a group G as in Proposition 5.17 cannot act freely, properly discontinuously by isometries on a Hadamard manifold of pinched negative curvature unless Q is virtually cyclic, as the normalizer $N_G(K_1)$ is not nilpotent.*

As an application of Proposition 5.15 we give the first examples of surface bundles over hyperbolic disks, with universal cover Gromov-hyperbolic. It has been an open question (cf. [KL08] [FM02]) to find purely pseudo Anosov surface groups in $MCG(S)$. The example below is a step towards this.

In [LS11] Leininger and Schleimer construct examples of disks (Q, d_Q) quasi-isometric to $\mathbb{H}^2$ and quasi-isometric embeddings $j : Q \to (Teich(S), d_T)$ such that $E \circ j : Q \to (Teich(S), d_e)$ is also a quasi-isometric embedding. Let $U(S, Q)$ be the pull-back bundle under j of the universal curve over $Teich(S)$ equipped with the natural path metric. By Proposition 5.15, $\widetilde{U(S, Q)}$ is a hyperbolic metric space, where $\widetilde{U(S, Q)}$ denotes the metric bundle obtained by taking fiberwise universal covers.

A brief sketch of Leininger-Schleimer's construction follows.

The curve complex $CC(S,x)$ of a surface with one puncture (or equivalently, a marked point x) admits a surjective map to $CC(S)$ such that the fiber over $\eta \in CC(S)$ is the Bass-Serre tree of the splitting of $\pi_1(S)$ over the cyclic groups represented by the simple closed curves in η. Suppose γ is a bi-infinite geodesic in $CC(S)$ coming from a geodesic in $Teich(S)$ lying in the thick part. Inside $CC(S,x)$ one has the space of trees over γ, and Leininger and Schleimer construct lines in each tree over γ whose union Q_1 is quasi-isometric to the hyperbolic plane. Using a branched cover-trick, Leininger and Schleimer construct from Q_1 a new disk $Q \subset CC(S')$ (for a closed surface S', which is a branched cover of S branched over the marked point of S) such that Q satisfies the hypotheses of Proposition 5.15.

References

[ABC⁺91] J. Alonso, T. Brady, D. Cooper, V. Ferlini, M. Lustig, M. Mihalik, M. Shapiro, and H. Short, *Notes on word hyperbolic groups*, Group Theory from a Geometrical Viewpoint (E. Ghys, A. Haefliger, A. Verjovsky eds.) (1991), 3–63.

[BF92] M. Bestvina and M. Feighn, *A Combination theorem for Negatively Curved Groups*, J. Differential Geom., vol 35 (1992), 85–101.

[BH99] M. Bridson and A Haefliger, *Metric spaces of nonpositive curvature*, Grundlehren der mathematischen Wissenchaften, Vol 319, Springer-Verlag (1999).

[Bow97] B. H. Bowditch, *Relatively hyperbolic groups*, preprint, Southampton (1997).

[Bow07] ———, *The Cannon-Thurston map for punctured surface groups*, Math. Z. 255 ((2007)), 35–76.

[CDP90] M. Coornaert, T. Delzant, and A. Papadopoulos, *Geometrie et theorie des groupes*, Lecture Notes in Math., vol.1441, Springer Verlag (1990).

[CT85] J. Cannon and W. P. Thurston, *Group Invariant Peano Curves*, preprint, Princeton (1985).

[CT07] ———, *Group Invariant Peano Curves*, Geom. Topol. 11 (2007), 1315–1355.

[Far98] B. Farb, *Relatively hyperbolic groups*, Geom. Funct. Anal. 8 (1998), 810–840.

[FM02] B. Farb and L. Mosher, *Convex cocompact subgroups of mapping class groups*, Geom. Topol. 6 (2002), 91–152.

[Gd90] E. Ghys and P. de la Harpe(eds.), *Sur les groupes hyperboliques d'apres Mikhael Gromov*, Progress in Math. vol 83, Birkhauser, Boston Ma. (1990).

[Gro85] M. Gromov, *Hyperbolic Groups*, in Essays in Group Theory, ed. Gersten, MSRI Publ.,vol.8, Springer Verlag (1985), 75–263.

[Gro93] ———, *Asymptotic Invariants of Infinite Groups*, in Geometric Group Theory,vol.2; Lond. Math. Soc. Lecture Notes 182, Cambridge University Press (1993).

[Ham05] U. Hamenstadt, *Word hyperbolic extensions of surface groups*, preprint, arXiv:math/0505244 (2005).

[Ham07] ———, *Geometry of complex of curves and teichmuller spaces*, in Handbook of Teichmuller Theory Vol. 1, EMS (2007), 447–467.

[Ham10] ———, *Stability of Quasigeodesics in Teichmuller Space*, Geom. Dedicata 146 (2010), 101–116.

[Kap08] M. Kapovich, *Problems on Boundaries of groups and Kleinian Groups*, http://www.aimath.org/pggt/Boundaries boundaries-version4.pdf (2008).

[KL08] R. P. KentIV and C. Leininger, *Shadows of mapping class groups: capturing convex cocompactness*, Geom. Funct. Anal. 18 (2008), 1270–1325.

[LMS08] C. Leininger, M. Mj, and S. Schleimer, *The universal Cannon–Thurston maps and the boundary of the curve complex* , to appear in Comment. Math. Helv., arXiv:0808.3521 (2008).

[LS11] C. Leininger and S. Schleimer, *Hyperbolic spaces in teichmuller spaces*, preprint (2011).

[Mit97] M. Mitra, *Ending Laminations for Hyperbolic Group Extensions*, Geom. Funct. Anal. 7 (1997), 379–402.

[Mit98a] ———, *Cannon-Thurston Maps for Hyperbolic Group Extensions*, Topology 37 (1998), 527–538.

[Mit98b] ———, *Cannon-Thurston Maps for Trees of Hyperbolic Metric Spaces*, J. Differential Geom. 48 (1998), 135–164.

[Mj06] M. Mj, *Cannon-Thurston Maps for Surface Groups*, preprint, arXiv:math.GT/0607509 (2006).

[Mj09] ———, *Mapping class groups and interpolating complexes: Rank*, J. Ramanujan Math. Soc. 24, no. 4 (2009), 341–357.

[Mj11] ———, *Cannon-Thurston Maps, i-bounded Geometry and a Theorem of McMullen*, Actes du séminaire Théorie spectrale et géométrie 28, Année 2009-10, arXiv:math.GT/0511104 (2011), 63–108.

[MM99] H. A. Masur and Y. N. Minsky, *Geometry of the complex of curves I: Hyperbolicity*, Invent. Math.138 (1999), 103–139.

[Mos96] L. Mosher, *Hyperbolic Extensions of Groups*, J. Pure Appl. Algebra 110(3) (1996), 305–314.

[Mos97] ———, *A hyperbolic-by-hyperbolic hyperbolic group*, Proc. Amer. Math. Soc. 125 (1997), 3447–3455.

[Mos03] ———, *Stable Teichmuller quasigeodesics and ending laminations* , Geom. Topol. 7 (2003), 33–90.

[MP11] M. Mj and A. Pal, *Relative Hyperbolicity, Trees of Spaces and Cannon-Thurston Maps*, arXiv:0708.3578, Geom. Dedicata 151(1) (2011), 59–78.

[MR08] M. Mj and L. Reeves, *A Combination Theorem for Strong Relative Hyperbolicity*, Geom. Topol. 12 (2008), 1777– 1798.

[Pal10] A. Pal, *Relatively Hyperbolic Extensions of Groups and Cannon-Thurston Maps*, Proc. Indian Acad. Sci. Math. Sci. 120 (1), arXiv:0801.0933 (2010), 57–68.

[Pal11] ———, *Notes on Stable Teichmuller quasigeodesics*, preprint, arXiv:1109.3794 (2011).

[Pap95] P. Papasoglu, *Strongly geodesically automatic groups are hyperbolic*, Invent. math. 121, (1995), 323– 334.

[Pap05] ———, *Quasi-isometry invariance of group splittings*, Ann. Math., vol. 161, no.2 (2005), 759–830.

RKM Vivekananda University, Belur Math, WB-711 202, India